



Tonian granitic magmatism of the Borborema Province, NE Brazil: A review



Ignez P. Guimarães^{a,*}, Maria de Fatima L. de Brito^b, Geysson de A. Lages^b,
Adejardo F. da Silva Filho^a, Lucilene Santos^a, Roberta G. Brasilino^b

^a Departamento de Geologia, Universidade Federal de Pernambuco, Recife 50740-530, Brazil

^b CPRM – Serviço Geológico do Brasil, Brazil

ARTICLE INFO

Article history:

Received 15 July 2015

Received in revised form

8 October 2015

Accepted 14 October 2015

Available online 21 October 2015

Keywords:

Tonian

Orthogneisses

Migmatites

Extension-related

Borborema Province

NE Brazil

ABSTRACT

Tonian granitoids, today augen-gneisses and migmatites, showing crystallization ages ranging from 870 Ma to 1000 Ma occur in the Borborema Province, NE Brazil. The majority of them have ages within the 925–970 Ma interval. Few intrusions with ages of ~1.0 Ga and <900 Ma occur in the Transversal and South subprovinces. The Tonian granitoids constitute the most expressive magmatic rocks of the Cariris Velhos event. The studied granitoids (herein CVG -Cariris Velhos granitoids) intrude slightly older bimodal (but mostly felsic) volcanic successions and metasedimentary sequences in the Transversal and South subprovinces. Tonian granitoids are unknown in the North subprovince. The CVG comprise mainly coarse-grained augen-gneisses of granite to granodiorite composition. Fe-rich biotite (annite) is the main mafic mineral phase, constituting up to 15% of the modal composition. Garnet, muscovite and tourmaline occur as accessory phases in many plutons.

The CVG augen-gneisses have high SiO₂ (>71%) and alkali contents, they vary from slightly peraluminous to slightly metaluminous, and from slightly magnesian to typical ferroan rocks. In the migmatized orthogneisses the SiO₂ contents are usually <70%. Trace element variations in the CVG are extensive, reflecting the migmatization recorded in some plutons and/or distinct sources. They are Ca-, Sr- and Nb-poor, showing variable Ba (100–1260 ppm), Rb (164–400 ppm) and Zr (144–408 ppm) contents, and high abundances of Y (>40 ppm). The chondrite normalized REE patterns are characterized by strong to moderate negative Eu anomalies (Eu/Eu* = 0.23–0.70). In general, the spidergram patterns show deep troughs at Ti, P, Ba and Sr and less pronounced Nb–Ta troughs. These patterns are similar to those reported for anorogenic granites evolved from mixtures of magmas from both crustal and mantle sources. The CVG exhibit T_{DM} model ages ranging from 1.9 to 1.1Ga, with slightly negative to slightly positive εNd(t) values, suggesting the involvement of distinct proportions of mantle and crustal components in the source of their protoliths.

There is no consensus in the literature about the tectonic setting of the CVG ie they have been related to either continental margin magmatic arc, with possible back-arc association, or extension-related setting, with generation of A-type granites. However, all the available geochemical data suggest that the CVG represent extension related magmatism. The geochemical signature associated to bimodal volcanism, including pyroclastic rocks, with similar ages, and absence, up to now, of evidence for metamorphism of Tonian age, support the hypothesis of extension - related magmatism.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

The Borborema Province (Almeida et al., 1981) constitutes a

region located in the northeast part of the South American continent, between the West Africa-São Luis craton to the north, and to the west by late Neoproterozoic (Brasiliano/Pan African) terranes that extend northward from central Brazil (Pimentel and Fuck, 1992; Pimentel et al., 1997, 2000) mostly covered by Phanerozoic sediments. According to Van Schmus et al. (2008), the Borborema Province resulted from the breakup of a Palaeoproterozoic

* Corresponding author. Universidade Federal de Pernambuco – Departamento de Geologia, CTG Cidade Universitaria, Recife, Pernambuco, Brazil.

E-mail address: ignez@ufpe.br (I.P. Guimarães).

supercontinent (Columbia: Rogers and Santosh, 2002, 2003; 2004) during the late Mesoproterozoic to early Neoproterozoic.

Van Schmus et al. (1995) proposed that the E–W trending Patos and Pernambuco shear zones divided the Borborema Province into three tectonic domains, later denominated subprovinces (Van Schmus et al., 2011): North, Transversal and South.

The North subprovince, located north of the Patos shear zone, comprises Paleoproterozoic basement, including some Archean nuclei, overlapped by Neoproterozoic metasedimentary sequences, and Ediacaran plutonic rocks. The South subprovince comprehends the region between the Pernambuco shear zone and the São Francisco craton, resulted from a tectonic collage of various blocks, terranes or domains with ages ranging from Archean to Neoproterozoic (Van Schmus et al., 2008).

The South subprovince comprises three domains (Fig. 1): (a) Sergipano, north of the São Francisco Craton; (b) Pernambuco–Alagoas (PEAL), north of the Sergipano domain, and (d) Riacho do Pontal. The Rio Preto belt is considered part of the South subprovince, but is not referred to as a domain by Van Schmus et al. (2011). The Sergipano domain or belt (Oliveira et al., 2010) is subdivided into five lithotectonic domains: Canindé, Poço Redondo–Marancó, Macururé, Vaza Barris, and Estância. The PEAL domain constitutes a region of high-grade gneisses, migmatites, and the largest Brazilian granitic intrusions in the Borborema Province, which lies to the north of the Sergipano domain. The Riacho do Pontal domain consists of metasedimentary and metavolcanic supracrustal sequences with late Neoproterozoic (Brasiliano) granites and some older gneisses.

The Transversal subprovince lies between the Patos and Pernambuco shear zones, within a major shear couple formed by dextral displacement of these two shear zones. This shear couple formed many faults, that caused fragmentation and clockwise block rotation, during the Pan African/Brasiliano Orogeny, which makes complex the reconstruction of the pre-Brasiliano geotectonic history of this domain. The Transversal subprovince was subdivided into two belts by Brito Neves (1983) and in four terranes by Santos (1995) and Santos et al. (1999). The terranes model has been contested by several works, and the Van Schmus et al. (2011) designation, will be used in this work.

Previous work (Brito Neves, 1983) did not consider the east branch of the Pernambuco shear zone as the limit of distinct domains, ie, the north part of the Pernambuco Alagoas domain continues in the Transversal subprovince of Van Schmus et al. (1995, 2008, 2011). Some recent works, carried out in the north part of the Pernambuco Alagoas domain and in the southeast part of the Transversal subprovince, have shown that the east branch of the Pernambuco shear zone is not a limit of distinct domains (Neves et al., 2012; Silva Filho et al., 2014a, b).

The Tonian (Cariris Velhos) suite comprises bimodal metavolcanic (mostly felsic) with small occurrence of pyroclastic, metasedimentary and metaplutonic rocks, yielding U–Pb zircon ages ranging from 870 Ma to 1000 Ma. This suite, mostly identified in the Transversal subprovince, is coeval with rocks from other parts of the South subprovince, notably in the Poço Redondo–Marancó subdomain of the Sergipano domain, in the Pernambuco Alagoas, Riacho do Pontal and Rio Preto domains.

At present, there is no integration for the Tonian orthogneisses dataset from distinct domains of the Borborema Province. The aim of this paper is to present a review of the Tonian orthogneisses from the Borborema Province, including geochronological, geochemical and Sm–Nd isotopic data from the literature and new geochemical and geochronological data. It is an attempt to constrain the Tonian orthogneisses geotectonic meaning in an integrated way, and doing so, to contribute to the understanding of the Cariris Velhos event in the Borborema Province.

2. Geological aspects of the Borborema Province

The Borborema Province consists of gneissic and migmatitic basement complexes, mostly formed during the Paleoproterozoic (1.98 Ga to 2.2 Ga), including minor Archean blocks, partially covered by Neoproterozoic metasedimentary and metavolcanic rocks (Van Schmus et al., 1995, 2008; Dantas et al., 1998; Fetter, 1999; Brito Neves et al., 2001; Kozuch, 2003; Guimarães et al., 2012).

Two contrasting tectonic models have been proposed to the late Neoproterozoic evolution of the Borborema Province: The accretionary model (Brito Neves and Cordani, 1991; Santos, 1995; Jardim de Sá, 1994 among others) in which the main shear zones represent boundaries of continental fragments or terranes amalgamated during the Pan-African – Brasiliano Orogeny. In opposition to the accretionary model, Neves (2003) and Neves et al. (2006) argued that the Borborema Province constitutes a consolidated block since 2.0 Ga ago (the single-block model).

Trompette (1997) suggested the continuation of the Transversal subprovince to the African side of West Gondwana, between the Adamaoua and Garoua shear zones in Cameroon (Median Shear Corridor). Some recent works (Silva Filho et al., 2014a; Neves et al., 2015) showed the at least in the east side of the Borborema Province, the Pernambuco lineament does not separate distinct blocks. However, there is a major contrast in Sm–Nd T_{DM} ages on either sides of the Patos shear zone: Archean model ages to the north and Mesoproterozoic model ages to the south. The distribution of the Sm–Nd T_{DM} model ages favors the hypothesis that the Patos shear zone represents a major terrane boundary (Brito Neves et al., 2001; Kozuch, 2003; Van Schmus et al., 2008). Geochemical and geochronological similarities between the Paleoproterozoic orthogneisses from the north and south of the Patos shear zone (Souza et al., 2007; Neves et al., 2006) suggest that they represent the basement of the supracrustal rocks with Mesoproterozoic T_{DM} model ages, recorded in the south of the Patos shear zone. They are now in the same level due to later vertical movement of the Patos shear zone. Dias et al. (2015), using seismic data, concluded that the Patos shear zone is likely to be a supracrustal structure, rather than a limit of lithospheric terranes, as suggested in the single-block model discussed above.

The Paleoproterozoic basement of the Borborema Province was affected by three major events, which took place at 2.35 Ga; 2.15 Ga and 2.0 Ga (Dantas, 1997; Neves et al., 2006, 2015; Souza et al., 2007). These events are interpreted as part of a Paleoproterozoic collage which led to the assembly of the Columbia Supercontinent of Rogers and Santosh (2002).

Sá et al. (2002); Accioly (2003) reported A-type metagranitoids and meta-anorthosites emplaced at 1.6 Ga and 1.77 Ga respectively, within the Transversal subprovince of the Borborema Province. In addition to the Paleoproterozoic orogenic cycle, the Borborema Province was affected by a Tonian event, the so-called Cariris Velhos event with magmatic ages within 870 Ma – 1000 Ma interval, and the Brasiliano (650–580 Ma) event.

The Cariris Velhos event in the Transversal subprovince is represented mainly by bimodal metavolcanic rocks, including pyroclastic varieties and granitic plutons, now orthogneisses (Santos, 1995; Brito Neves et al., 2001; Kozuch, 2003; Guimarães et al., 2012) and, restricted occurrence of metasediments (muscovite-biotite gneisses, garnet-biotite schists, marbles and BIF) intercalated with metavolcanic rocks. Jardim de Sá et al. (1988), reported the first Tonian age in the Borborema Province. Using Rb–Sr whole rock, they defined an isochronous age of 968 ± 35 Ma, for the orthogneisses of the Afeição suite in the South subprovince (Riacho do Pontal domain) of the Borborema Province, which has been interpreted as representing mixed ages. Later, Van Schmus et al.




Fig. 1. a) The subprovinces of the Borborema Province, according to Van Schmus et al. (2011), showing the areas detailed in figures b, c and d. Sketch geological maps of: b) the Transversal and part of the South subprovinces (part of the Pernambuco – Alagoas domain), c) the Pernambuco–Alagoas and Sergipano domains, and d) Riacho do Pontal and Rio Preto domains showing the location of plutons with available U–Pb zircon geochronological data. 1- (925 ± 10 Ma – Jardim), 2- (952 ± 7 Ma) and 3- (942 ± 22 Ma – Fazenda Agra) are from Brito Neves et al. (2001). 4 – Sítio do Icó (1005 ± 5 Ma – this work). 5 – (960 ± 4 Ma Morro do Cruzeiro Pluton) from Guimarães et al. (2012). 6 – (958 ± 9 Ma, Ambó pluton), 7 – (926 ± 12 Ma), 8 – (941 ± 17 Ma – Imaculada pluton), 9- (959 ± 5 Ma – Queimadas orthogneisses), 10 – (963 ± 14 Ma- Serra Talhada orthogneisses) all from Kozuch (2003), 11- (Tabira pluton – 972 ± 5 Ma) from Leite et al. (2000). 12 – (999 ± 50 Ma –Canoas gneiss) from Van Schmus et al., 1995). 13 – (Recanto – 1037 ± 30 Ma) from Santos (1995). 14 – Riacho do Forno (993 ± 5 Ma – this work), 15 – (Lobo – 974 ± 6 Ma) from Cruz et al. (2014), 15* – (994 ± 25 Ma – Lobo pluton) from Brito and Freitas (2011). 16- (956 ± 2 Ma – Rocinha pluton) from Cruz et al. (2014), 17 – (870 ± 8 Ma – Pinhões pluton) from Neves et al. (2012), 18 – (960 ± 11 Ma – Riacho do Tigre metavolcanics) of Accioly et al. (2010). c) 19 – (947 ± 6 Ma – Flores) from Brito et al. (2008), 20 – (972 ± 30 Ma – sample SI-100) from Silva Filho et al. (2014a, b), 21 – (SI-125 – 983 ± 9 Ma – this Work), 22 – (979 ± 3 Ma – Poço Redondo migmatites) and 23 – (952 ± 2 Ma – Serra Negra pluton) are from Carvalho (2005). d) 24 – (966 ± 5 Ma – Afeição augen gneisses) and 24* – (1001 ± 4 Ma – granite sill) are from Caxito et al. (2014) and 24 – (969 ± 11 Ma – Serra da Pintada) from Aquino and Batista (2011). The Sketch geological maps are modified from Medeiros (2004).

(1995), using U–Pb zircon TIMS data in a sample from the same suite, obtained similar age (966 ± 10 Ma), and interpreted it as the crystallization age of the orthogneisses protolith.

There is no consensus in the literature concerning the tectonic setting of the Tonian granitoids. They are related to either continental margin magmatic arc, with possible back–arc association

(Brito Neves et al., 2000; Santos et al., 2010; Oliveira et al., 2010; Kozuch, 2003; Caxito et al., 2014, Van Schmus et al., 2011), or extension-related setting, with generation of A-type granites (Oliveira et al., 2010; Guimarães et al., 2012; Cruz et al., 2014; Aquino and Batista, 2011).

The Pan African/Braziliano event (650 Ma – 550 Ma) affected the

entire Borborema Province and was responsible for low-to high-temperature and low-pressure metamorphism, a great abundance of granitic intrusions, and the development of continental-scale transcurrent shear zones. The Brasiliano event was also responsible by the actual structural framework of the Borborema Province.

3. Geology and petrographic aspects of the Cariris Velhos granitoids

3.1. Transversal subprovince

The Cariris Velhos orthogneisses encompass a 50–100 km wide belt that extends for more than 700 km west–southwestwards from the Atlantic coastline. The orthogneisses, representing granitic plutons, intrude a metavolcano - metasedimentary sequence, which crops out on both sides of the gneissic core. The metasediments with Tonian deposition ages appear to have a restrict distribution (Guimarães et al., 2012), being recognized only at east of the Serra do Caboclo shear zone, the so-called metavolcano–metasedimentary Riacho Gravatá Complex (Bittar, 1998). However, late Neoproterozoic sediments have received contribution from the Cariris Velhos suite and were deposited within the same belt.

According to Bittar (1998) the Riacho Gravatá Complex comprises three stratigraphic units: Serra do Mocambo (NNE), Macacos-Piauí (center) and Serra da Pinheira (SW). U–Pb zircon TIMS (Kozuch, 2003) and SHRIMP (Guimarães et al., 2012) data show that the metavolcanic rocks from the three stratigraphic units proposed by Bittar (1998) yield ages ranging from 970 ± 15 Ma to 986 ± 19 Ma.

The Serra do Caboclo shear zone is the boundary between the Riacho Gravatá Complex and the Pianco-Alto Brigida domain. The Riacho Gravatá Complex is bounded to the east by the Teixeira-Terra Nova high. Bittar (1998) described the Riacho Gravatá Complex as a metavolcano-metasedimentary sequence metamorphosed under low to medium metamorphic grade; it includes muscovite \pm graphite biotite schists, metagreywackes, marble, metatuffs, metaultramafic rocks, metasiltite, bimodal (but mostly felsic) metavolcanic rocks, and local occurrences of banded iron formation. The metavolcanic mafic rocks are greenschist and amphibolite grade.

The studied orthogneisses represent a set of individual small plutons (<80 km²) comprising mainly coarse-grained augen-gneisses of granite to granodiorite composition. Fe-rich biotite (annite) is the main mafic mineral phase constituting up to 15% of the modal composition. Garnet, muscovite and tourmaline occur in many plutons. Zircon, apatite and allanite are the main accessory phases. Some plutons comprise strongly migmatized gneisses and some others are composed of banded orthogneisses. However, most consist of augen-gneisses, and some of them (Ambó and Morro do Cruzeiro plutons among others) show enclaves preserved from the Brasiliano deformation.

Fine grained migmatitic orthogneisses have been described in the east part (Lages et al., 2014) and in the south part of the Cariris Velhos belt. The migmatites from east part comprise amphibole-rich layers intercalated with felsic layers, folded and migmatized. In the south, the migmatites are also fine grained and have biotite but smaller modal amount of amphiboles. The other Tonian orthogneisses of the Borborema Province have only biotite as the main mineral mafic phase.

3.2. South subprovince

3.2.1. Sergipano domain

The Sergipano domain has a triangular shape, with a NW–SE

structural strike (Fig. 1). It represents the southern segment of a Brasiliano continental collision zone which continues into Central Africa as the North Equatorial fold belt and may have resulted from the closure of a large intracontinental basin, along the northern margin of the São Francisco-Congo Craton (Davison and Souza, 1989). According to Trompette (1997, 2000) the Sergipano domain comprises the western segment of the Oubangide orogen, which extends into NW Africa in pre-drift reconstructions.

Davison and Souza (1989) recognized five lithotectonic domains in the Sergipano domain, namely from N to S: Canindé, Poço Redondo-Marancó, Macururé, Vaza Barris and Estância, which are mutually separated by Neoproterozoic shear zones. The Poço Redondo-Marancó domain is composed of two units: Poço Redondo, which constitutes the basement, and the Marancó supracrustal rocks (Oliveira et al., 2010).

Carvalho (2005) and Oliveira et al. (2010) described occurrences of late Mesoproterozoic to early Neoproterozoic migmatites (Poço Redondo) and augen gneisses (Serra Negra), which constitute the basement of the Marancó – Poço Redondo domain. The Serra Negra augen – gneisses occur as an elongated WNW – ESE oriented pluton, in the south part of the Poço Redondo subdomain. The gneisses show tectonic contacts, made mainly through transcurrent shear zones, with the Poço Redondo migmatites (Carvalho, 2005). Metasedimentary xenoliths are common within the Serra Negra orthogneisses.

The Poço Redondo migmatites constitute a belt that extends in the WNW–ESE direction, for more than 30 km long and 8 km wide. Silva Filho et al. (1997) firstly described these rocks as homogeneous and heterogeneous migmatites. They suggest that part of these migmatites has a metasedimentary source.

The timing of deformation was constrained by Oliveira et al. (2010), using ⁴⁰Ar–³⁹Ar dating of amphiboles, at 625 ± 3 Ma and muscovite plates from supracrustal rocks at 612 ± 7 Ma. The data support that the deformation has a late Neoproterozoic (Pan African/Brasiliano) age.

3.2.2. Riacho do Pontal domain

The Riacho do Pontal domain is located just south of the westernmost branch of the Pernambuco shear zone (Fig. 1a). According to Van Schmus et al. (2011) the Riacho do Pontal domain may represent the SW continuation of some of the terranes in the Transversal subprovince, including the Cariris Velhos belt.

Oliveira (1998) and Caxito et al. (2014) subdivided the Riacho do Pontal domain into three tectono-stratigraphic zones or domains: external, central and internal zones.

The external zone comprehends a south-verging nappe system in the southernmost part of the domain. It comprises a supracrustal sequence (Casa Nova Group) composed mainly of clastic metasedimentary rocks such as quartzite, garnet-mica schist containing staurolite and kyanite, greywacke and marble intercalations. According to Caxito et al. (2012), this supracrustal sequence is related to a syn-orogenic basin.

The central zone comprises metamafic rocks with MORB signature (Moraes, 1992), intercalated within a metasedimentary sequences of metacherts, metagreywackes and garnet mica schist (Monte Orebe Complex). Angelim (1988) based in field relationships suggested a Tonian age for the Monte Orebe Complex. According to Caxito et al. (2014), the central zone marks an ancient suture zone between two distinct lithospheric blocks: the São Francisco Craton to the south, without Tonian-related intrusions, and the internal zone block, which contains the Afeição suite. However, few U–Pb zircon SHRIMP data, presented by Brito Neves et al. (2015), suggest a Neoproterozoic deposition age, between 650 Ma and 850 Ma, for part of the Monte Orebe sediments.

The internal zone comprehends granitic plutonism, today augen




Fig. 2. a) Zircon SEM – CL pictures for sample GL-544 (Sítio do Icó). b) Concordia diagram for sample GL-544 (location 4 in Fig. 1b).

gneisses of the Afeição suite, reworked migmatitic basement and predominantly metasedimentary sequences, with minor volcanic contribution (Gomes and Vasconcelos, 1991; Angelim and Kosin, 2001).

The supracrustal sequences of the Riacho do Pontal domain are intruded by plutons with crystallization ages, mostly Rb – Sr data, within the 668 – 550 Ma interval (Jardim de Sá et al., 1988, 1992; Jardim de Sá, 1996; Caxito et al., 2014).

3.2.3. Rio Preto domain

The Rio Preto domain borders the northwestern São Francisco Craton, southwest of the Riacho do Pontal domain (Fig. 1a). According to Caxito et al. (2014), it represents the geological link between the Borborema and the Northern Tocantins orogenic provinces of Brazil. The Rio Preto domain comprises a Paleoproterozoic basement, extensively reworked during the Brasiliano Orogeny (Egydio-Silva et al., 1989) and two metasedimentary units: 1) Canabrinha Formation to the south, which is composed of a para-derived sequence constituted by conglomerates and sediments similar to turbidites (Egydio-Silva et al., 1989) with Neoproterozoic (~900–600 Ma) deposition ages (Caxito et al., 2014). 2) The Formosa Formation to the north, comprises fine-grained sediments, associated with cherts and probably submarine volcanic activity, now represented by greenschists (Caxito et al., 2012). The deposition of the Formosa Formation sediments appear to have occurred during the Orosirian Period (~1.9 Ga – Caxito et al., 2014).

The Serra da Pintada pluton comprises leucocratic coarse – grained augen-gneisses of syenogranite composition (Aquino and Batista, 2011). The pluton forms an elongated body, showing maximum axis with E–W direction, intruded into Paleoproterozoic

orthogneisses. Three more plutons (Algodões granitoids) with Tonian ages are described in the Rio Preto domain (Aquino, 2014), less than 15 km southwest of the Serra da Pintada pluton. The plutons comprise coarse – to medium - grained augen gneiss, locally mylonitic, with composition ranging from alkali-feldspar granite to syenogranite.

4. Geochronological data

4.1. Transversal subprovince

This subprovince has the majority of the geochronological data available for Tonian orthogneisses. A large number of U–Pb zircon ages, for Tonian orthogneisses from the Cariris Velhos belt of the Transversal subprovince, was obtained by Van Schmus et al. (1995) and Kozuch (2003), using standard isotope-dilution, thermalization mass spectrometry (ID-TIMS) methods. This dataset was important to constraint the crystallization ages of the orthogneisses protolith. The U–Pb zircon TIMS ages came out very similar to that defined by Brito Neves et al. (2000, 2001) using whole rock Rb–Sr methodology. Guimarães et al. (2012) provided U–Pb zircon SHRIMP age of 960 ± 4 Ma for one orthogneisse sample of the Cariris Velhos belt (Morro do Cruzeiro pluton - Fig. 1b). This pluton is constituted by coarse – grained augen - gneisses having granitic compositions, similar to that of the granitoids from the Ambó pluton which has U–Pb zircon TIMS crystallization age of 958 ± 9 Ma (Kozuch, 2003). Twenty-five zircon grains were analyzed and all analyses are concordant and no significantly older inherited cores were recorded. Guimarães et al. (2012) also reported very narrow zircon overgrowths, and in attempt to define their ages, seven overgrowths were analyzed. However, the obtained $^{206}\text{Pb}/^{238}\text{U}$ ages (678 Ma to 881 Ma) were not able to define the metamorphic age.

Besides the Ambó pluton, Kozuch (2003) analyzed four more plutons from the same belt (Fig. 1b), by U–Pb zircon TIMS, and obtained the following ages: Imaculada gneisses - 941 ± 17 Ma; Fazenda Arroz orthogneisses – 926 ± 12 Ma; Queimadas gneisses – 959 ± 5 Ma and Serra Talhada gneisses 963 ± 28 Ma. In the Serra Talhada gneisses, Kozuch (2003) recorded a younger zircon population that gave an age of 626 ± 28 Ma, which was interpreted as the Brasiliano metamorphic age. This age is similar to Ar–Ar ages in amphiboles obtained by Oliveira et al. (2010) in the Sergipano domain. Leite et al. (2000) reported U–Pb zircon TIMS age of 969 ± 4 Ma for an augen - gneiss sample (Fig. 1b), which they called Tabira pluton. In the same sample, Leite et al. (2000) also analyzed one fraction of titanite, which gave a concordant age of 612 ± 9 Ma, interpreted as a metamorphic age. Guimarães et al., 2012 reported granitoids, denominated as Tabira pluton, intruded into the orthogneisses described by Leite et al. (2000) and has U–Pb zircon SHRIMP crystallization age of 593 ± 7 Ma (Guimarães et al., 2012), which is similar to the U–Pb titanite age recorded by Leite et al. (2000).

Felsic metavolcanic rocks from the Cariris Velhos belt, within the so-called Riacho Gravatá Complex, have U–Pb TIMS age ranging from 913 ± 25 Ma to 980 ± 10 Ma (Kozuch, 2003) and U–Pb zircon SHRIMP ages of 996 ± 13 Ma and 986 ± 10 Ma (Guimarães et al., 2012). Brito Neves et al. (2001) reported U–Pb zircon TIMS age of 925 ± 10 Ma and 952 ± 7 Ma (Jardim pluton) in granitoids from the east part of the Cariris Velhos belt (Fig. 1b). Brito Neves et al. (2001) also suggest that the metasediments in this specific area have Tonian ages. However, U–Pb zircon SHRIMP data presented by Guimarães et al. (2012) show that the deposition of these sediments occurred during the Pan African/Brasiliano orogeny.

Thirty six zircon grains were extracted from a fine-grained, amphibole rich layer of a migmatized orthogneisse sample from the east part of the Cariris Velhos belt (Sítio do Icó – Fig. 1b). U–Pb

Table 1
Summary of U/Pb SHRIMP data for sample GL-544 – Sítio do Icó.

Samples	%	ppm		Ratios						ρ	Apparent ages						Conc.(%)		
		Comm	U	Th	Th/U	206 Pb		207 Pb			$\pm 1\sigma$		207 Pb (Ma)		207 Pb (Ma)			206 Pb (Ma)	
						204 Pb	206 Pb	235U	238U		206 Pb	$\pm 1\sigma$	206 Pb	235U	238U				
						206													
016-Z9			0.68	59225	0.072616	0.46	1.534	0.75	0.153176	0.60	0.75	1003	9	944	5	919	5.14	92	
045-Z27			0.81	82962	0.073030	0.57	1.542	0.89	0.153183	0.68	0.73	1015	12	947	5	919	5.84	91	
015-Z8			0.48	251120	0.072850	0.48	1.581	0.75	0.157374	0.58	0.71	1010	10	963	5	942	5.04	93	
046-Z28			0.87	127713	0.072716	0.51	1.582	0.80	0.157797	0.62	0.72	1006	10	963	5	945	5.41	94	
041-Z25			0.87	174645	0.072122	0.50	1.595	0.80	0.160402	0.62	0.73	989	10	968	5	959	5.50	97	
005-Z2			0.53	84051	0.071346	0.83	1.601	1.30	0.162749	1.00	0.75	967	17	971	8	972	9.05	100	
048-Z30			0.91	109495	0.073855	1.26	1.602	1.41	0.157320	0.63	0.64	1038	25	971	9	942	5.55	91	
040-Z24			0.44	186668	0.071880	0.34	1.607	0.67	0.162136	0.58	0.82	983	7	973	4	969	5.21	99	
024-Z15			0.82	87259	0.072298	0.74	1.617	0.91	0.162221	0.53	0.71	994	15	977	6	969	4.79	97	
042-Z26			0.51	138735	0.073084	0.88	1.622	1.06	0.160962	0.59	0.71	1016	18	979	7	962	5.26	95	
047-Z29			0.51	315027	0.073021	0.45	1.626	0.80	0.161531	0.66	0.79	1015	9	980	5	965	5.89	95	
027-Z16			0.86	188356	0.072836	0.47	1.630	0.69	0.162311	0.50	0.63	1009	10	982	4	970	4.49	96	
039-Z23			0.93	191008	0.072572	0.39	1.644	0.67	0.164270	0.54	0.75	1002	8	987	4	980	4.96	98	
034-Z20			0.77	90941	0.072554	0.44	1.651	0.80	0.165021	0.67	0.80	1002	9	990	5	985	6.13	98	
021-Z12			0.94	68449	0.072644	0.33	1.657	0.56	0.165459	0.45	0.69	1004	7	992	4	987	4.09	98	
036-Z22			0.73	73451	0.074654	1.18	1.665	1.37	0.161798	0.70	0.71	1059	24	995	9	967	6.25	91	
018-Z11			0.40	114661	0.074410	0.95	1.674	1.07	0.163191	0.50	0.60	1053	19	999	7	974	4.51	93	
033-Z19			0.79	175713	0.072370	0.42	1.679	0.76	0.168282	0.64	0.80	996	9	1001	5	1003	5.90	101	
022-Z13			0.55	89653	0.073025	0.44	1.697	0.76	0.168513	0.62	0.78	1015	9	1007	5	1004	5.78	99	
035-Z21			0.29	69472	0.073201	0.43	1.697	0.82	0.168113	0.70	0.83	1020	9	1007	5	1002	6.52	98	
009-Z4			0.63	56826	0.074494	0.57	1.711	0.77	0.166569	0.53	0.60	1055	11	1013	5	993	4.87	94	
029-Z18			0.51	145602	0.073026	0.40	1.713	0.64	0.170149	0.50	0.70	1015	8	1013	4	1013	4.70	100	
017-Z10			0.54	71993	0.076295	0.61	1.721	0.86	0.163564	0.61	0.65	1103	12	1016	6	977	5.53	89	
004-Z1			0.71	117145	0.072804	0.57	1.730	0.84	0.172386	0.63	0.69	1009	11	1020	5	1025	5.94	102	
006-Z3			0.92	56317	0.074142	0.74	1.740	0.91	0.170201	0.54	0.71	1045	15	1023	6	1013	5.07	97	
023-Z14			0.90	41067	0.075086	0.59	1.763	0.85	0.170280	0.61	0.66	1071	12	1032	5	1014	5.72	95	
010-Z5			0.67	20429	0.075809	0.55	1.768	0.83	0.169193	0.62	0.69	1090	11	1034	5	1008	5.76	92	
011-Z6			0.53	4034	0.081642	0.54	1.781	0.81	0.158219	0.60	0.68	1237	11	1039	5	947	5.26	77	
012-Z7			0.43	22041	0.079917	1.09	1.791	1.37	0.162508	0.84	0.78	1195	21	1042	9	971	7.61	81	
028-Z17			0.80	5609	0.077710	0.54	1.803	0.79	0.168254	0.58	0.66	1139	11	1046	5	1002	5.34	88	

zircon analysis were carried out with a Thermo-Finnigan multi-collector LA-ICP-MS at the Geochronology Laboratory of Brasília University. An apparent age of 1005 ± 5 Ma (Fig. 2a, b; Table 1) was obtained in the upper intercept. This age is interpreted as the crystallization age of the gneiss protolith. The zircon grains are euhedral to sub-euhedral and zoned; overgrowths are rare and, when they occur, they are very narrow (Fig. 2a). Xenocrystic cores analyzed from 3 grains; yielded $^{206}\text{Pb}/^{238}\text{U}$ ages ranging from 1013 ± 5 Ma to 1025 ± 6 Ma. The $^{232}\text{Th}/^{238}\text{U}$ ratios range from 0.40 to 0.93, which are compatible with magmatic source. No substantially older cores were found. The zircon grain shapes (Fig. 2b) are similar to those recorded by Guimarães et al. (2012) in a metavolcanic rock from the Riacho Gravatá Complex.

A sample from a migmatized orthogneiss (Sample AM-536), of muscovite monzogranite composition, was collected from the Riacho do Forno type locality (Fig. 1b). Thirteen spots were analyzed from 12 zircon grains. U–Pb analysis were done by SHRIMP at the Geochronological Research Center (CPGeo) of the Geosciences Institute of the São Paulo University, Brazil. Details of the analytical procedures and calibration of the device are described by Stern (1997), Williams (1998) and Sato et al. (2008).

Zircon grains are prismatic with length/width ratios up to 1 to 4 (Fig. 3a), however, rounded shape grains were also recorded. Oscillatory zoning is ubiquitous. Overgrowths are not apparent, but many outer zones have embayment (Fig. 3a). Six substantially older cores were found (Table 2), showing $^{206}\text{Pb}/^{238}\text{U}$ ages ranging from 1777 ± 45 Ma to 2333 ± 60 Ma. A Concordia age was defined at 993 ± 9 Ma (Fig. 3b). This age is interpreted as the crystallization age of the orthogneiss protolith. This age helps to improve the U–Pb zircon TIMS age of 1037 ± 30 Ma obtained by Santos (1995) for the Recanto – Riacho do Forno orthogneisses.

The ages of Serra do Icó and Recanto – Riacho do Forno

orthogneisses are slightly older than those of the other orthogneisses from the Cariris Velhos belt, but they are similar to those recorded in felsic metavolcanic rocks of the Riacho Gravatá Complex (Guimarães et al., 2012). One sample, analyzed from the Canoa orthogneiss, from the south part of the Cariris Velhos belt (Fig. 1b), shows age of 999 ± 50 Ma (Van Schmus et al., 1995). The error is too large and more data is necessary to better constraint the crystallization age of the gneiss protolith.

Outside the main belt, Tonian LA-ICP-MS zircon ages are reported in metavolcanic acid rock from the Riacho do Tigre Complex (Fig. 1b), with crystallization age of 960 ± 11 Ma (Accioly et al., 2010). Metaplutonic rocks of younger Tonian ages, the Pinhões orthogneisses (870 ± 8 Ma – Fig. 1b) are recognized in the east part of the Transversal subprovince (Rio Capibaribe domain) by Neves et al. (2012), and may represent the end of the Tonian event in the Borborema Province.

4.2. The south subprovince

4.2.1. Pernambuco Alagoas (PEAL) domain

Geochronological data are available for the west part of the PEAL domain, in three granitic plutons intruded just south of the Pernambuco shear zone (Fig. 1c). The granitoids have ages of 994 ± 25 Ma (Lobo 1 pluton – Brito and Marinho, 2014; Brito and Freitas, 2011); 974 ± 8 Ma (Lobo 2- pluton – Cruz et al., 2014) and 956 ± 2 Ma (Cruz and Accioly, 2013).

Within the east part of the PEAL domain, Brito et al. (2008) reported age of 947 ± 6 Ma for the Flores pluton (Fig. 1c) which has a maximum N–S oriented axis. U–Pb zircon SHRIMP data from a metatexite (sample SI-100, Fig. 1c), west of the Flores orthogneisses gave an age of 972 ± 30 Ma (Silva Filho et al., 2014b). Both ages are interpreted as the protoliths crystallization ages.




Fig. 3. a) Zircon SEM – CL pictures for sample AM-536 (Riacho do Forno – location 14 in Fig. 1b). b) Concordia diagram for sample AM-536.

Zircon grains were extracted from sample SI-125, collected from a migmatite close to the sample SI-100 (Fig. 1c). Twenty zircons grains were analyzed at SHRIMP RG (Sensitive High Resolution Ion

Microprobe) from the Research School of Earth Sciences of the Australia National University. Analytical methods and data reduction followed those detailed by Williams (1998) and Stern (1997) using the FC1 standard (1099 Ma; Paces and Miller, 1993).

The majority of the zircon grains are euhedral to subhedral and show igneous zoning. Narrow overgrowths occur in many grains. (Fig. 4a). Some grains exhibit corroded rims. No substantially older cores were found.

In a Concordia diagram (Fig. 4b) a mean $^{207}\text{Pb}/^{206}\text{Pb}$ age of $983 \pm 9\text{Ma}$ (MSWD = 0.78, probability of fit = 0.62) was defined for nine cores of the zircon grains (Fig. 4, Table 3), and interpreted as the crystallization age of the migmatite protoliths. A mean $^{207}\text{Pb}/^{206}\text{Pb}$ age of $664 \pm 8\text{Ma}$ (MSWD = 0.88, probability of fit = 0.50) was defined for seven zircon rims and interpreted as a metamorphic age.

Oxygen zircon data (Silva Filho et al., 2014a, b) from sample SI-100 show $\delta^{18}\text{O}$ values ranging from +3.45‰ to +5.53‰, consistent with mantle derived magmas, crystallized in a closed system. In spite of few Tonian granitoids recognized in the Pernambuco Alagoas domain up to now, many detrital zircon grains with Tonian ages are reported in metasediments of the PEAL domain (Neves et al., 2015; Silva Filho et al., 2014a, b).

4.2.2. Sergipano domain

The occurrences of granitoids with Tonian ages within the Sergipano domain are restricted to the Poço Redondo - Marancó subdomain (Fig. 1c). The available dataset comprises ages of the Serra Negra augen - gneisses and Poço Redondo migmatite. Carvalho (2005) reported two U–Pb SHRIMP ages of $980 \pm 4\text{Ma}$ and $961 \pm 32\text{Ma}$ for two samples of the Poço Redondo paleosome migmatites and an age of $951 \pm 2\text{Ma}$ to the Serra Negra augen-gneisses. The ages are interpreted as the crystallization ages of the migmatite and augen-gneisses protoliths. A zircon population with Concordia age of $967 \pm 2\text{Ma}$, recorded in the Serra Negra augen-gneisses, is interpreted as inherited from the Poço Redondo migmatite country rock (Carvalho, 2005).

The Marancó metasediments have a large population of zircon grains with ages within the 980–1000 Ma interval, few Paleoproterozoic and Archean ages and some ages within the 960–970

Table 2 - Summary of U–Pb SHRIMP zircon data for sample SI-125, from the PEAL domain.

Grain/Spot	% $^{206}\text{Pb}_c$	ppm U	ppm Th	$^{232}\text{Th}/^{238}\text{U}$	±%	(1) ppm $^{206}\text{Pb}^*$	(1) $^{206}\text{Pb}/^{238}\text{U}$ Age	(1) $^{207}\text{Pb}/^{206}\text{Pb}$ Age	% Disc.	(1) $^{207}\text{Pb}^*/^{206}\text{Pb}^*$	±%	(1) $^{207}\text{Pb}^*/^{235}\text{U}$	±%	(1) $^{206}\text{Pb}^*/^{238}\text{U}$	±%	err corr
1	0.07	187	117	0.65	0.26	25	952 ±12	972 ±15	+2	0.0715	0.74	1.569	1.5	0.1591	1.3	0.87
2.1	0.49	1603	155	0.10	7.26	137	611 ±8	669 ±21	+9	0.0618	0.96	0.848	1.6	0.0995	1.3	0.80
3.1	0.01	273	195	0.74	0.53	39	988 ±17	989 ±10	+0	0.0721	0.50	1.647	1.9	0.1656	1.9	0.97
4.1	0.20	526	3	0.01	1.07	45	617 ±16	677 ±16	+9	0.0621	0.73	0.861	2.8	0.1005	2.7	0.96
5.1	0.17	168	100	0.61	0.29	22	933 ±11	967 ±18	+4	0.0713	0.88	1.532	1.5	0.1558	1.3	0.82
6.1	0.02	141	58	0.42	0.32	20	973 ±14	982 ±16	+1	0.0719	0.78	1.615	1.8	0.1630	1.6	0.90
7.1	–	655	5	0.01	0.80	58	636 ±6	679 ±10	+7	0.0621	0.46	0.889	1.2	0.1038	1.1	0.92
8.1	0.17	197	70	0.37	2.43	24	852 ±10	898 ±17	+6	0.0690	0.83	1.344	1.5	0.1413	1.2	0.83
9.1	0.08	903	7	0.01	0.71	79	621 ±6	666 ±10	+7	0.0618	0.48	0.862	1.2	0.1012	1.1	0.91
10.1	0.04	200	91	0.47	6.46	24	836 ±22	914 ±15	+9	0.0695	0.71	1.326	2.9	0.1384	2.8	0.97
11.1	–	236	181	0.79	1.19	32	950 ±17	974 ±12	+3	0.0716	0.58	1.568	2.0	0.1588	1.9	0.96
12.1	0.01	752	5	0.01	0.92	68	648 ±9	660 ±9	+2	0.0616	0.43	0.899	1.6	0.1058	1.5	0.96
13.1	0.26	548	129	0.24	4.36	55	713 ±15	775 ±28	+8	0.0650	1.33	1.048	2.6	0.1170	2.2	0.86
14.1	0.06	197	115	0.60	0.45	27	965 ±17	983 ±15	+2	0.0719	0.72	1.601	2.1	0.1615	1.9	0.94
15.1	0.08	195	120	0.64	0.25	27	972 ±19	1009 ±15	+4	0.0728	0.73	1.634	2.2	0.1627	2.1	0.95
16.1	0.05	90	59	0.68	1.23	12	952 ±10	998 ±20	+5	0.0724	1.00	1.589	1.5	0.1591	1.2	0.76
17.1	0.41	215	85	0.41	1.63	21	682 ±7	961 ±24	+31	0.0711	1.20	1.095	1.6	0.1116	1.1	0.68
18.1	0.18	258	181	0.73	0.26	33	891 ±16	975 ±16	+9	0.0716	0.77	1.463	2.0	0.1482	1.9	0.93
22.1	0.04	869	8	0.01	0.69	78	640 ±8	660 ±10	+3	0.0616	0.44	0.887	1.4	0.1045	1.4	0.95
21.1	0.03	717	4	0.01	0.95	64	642 ±9	650 ±11	+1	0.0613	0.49	0.885	1.6	0.1048	1.5	0.95

Errors are 1-sigma; Pb_c and Pb^* indicate the common and radiogenic portions, respectively. Error in Standard calibration was 0.25% (not included in above errors but required when comparing data from different mounts) 0.1 – Common Pb corrected using measured ^{204}Pb . 2-Common Pb corrected by assuming $^{206}\text{Pb}/^{238}\text{U}$ – $^{207}\text{Pb}/^{235}\text{U}$ age-concordance.




Fig. 4. a) Cathodoluminescence images showing the internal and external characteristics of zircon grains extracted from sample SI-125 and b) U–Pb Concordia diagram of SHRIMP analyses of zircon from sample SI-125.

Table 3

Summary of U/Pb SHRIMP data for sample SI-125.

Grain/Spot	% $^{206}\text{Pb}_c$	ppm U	ppm Th	$^{232}\text{Th}/^{238}\text{U}$	±%	(1) ppm $^{206}\text{Pb}^*$	(1) $^{206}\text{Pb}/^{238}\text{U}$ Age	(1) $^{207}\text{Pb}/^{206}\text{Pb}$ Age	% Disc.	(1) $^{207}\text{Pb}^*/^{206}\text{Pb}^*$	±%	(1) $^{207}\text{Pb}^*/^{235}\text{U}$	±%	(1) $^{206}\text{Pb}^*/^{238}\text{U}$	±%	err Cor
1	0.07	187	117	0.65	0.26	25	952 ±12	972 ±15	+2	0.0715	0.74	1.569	1.5	0.1591	1.3	0.87
2.1	0.49	1603	155	0.10	7.26	137	611 ±8	669 ±21	+9	0.0618	0.96	0.848	1.6	0.0995	1.3	0.80
3.1	0.01	273	195	0.74	0.53	39	988 ±17	989 ±10	+0	0.0721	0.50	1.647	1.9	0.1656	1.9	0.97
4.1	0.20	526	3	0.01	1.07	45	617 ±16	677 ±16	+9	0.0621	0.73	0.861	2.8	0.1005	2.7	0.96
5.1	0.17	168	100	0.61	0.29	22	933 ±11	967 ±18	+4	0.0713	0.88	1.532	1.5	0.1558	1.3	0.82
6.1	0.02	141	58	0.42	0.32	20	973 ±14	982 ±16	+1	0.0719	0.78	1.615	1.8	0.1630	1.6	0.90
7.1	–	655	5	0.01	0.80	58	636 ±6	679 ±10	+7	0.0621	0.46	0.889	1.2	0.1038	1.1	0.92
8.1	0.17	197	70	0.37	2.43	24	852 ±10	898 ±17	+6	0.0690	0.83	1.344	1.5	0.1413	1.2	0.83
9.1	0.08	903	7	0.01	0.71	79	621 ±6	666 ±10	+7	0.0618	0.48	0.862	1.2	0.1012	1.1	0.91
10.1	0.04	200	91	0.47	6.46	24	836 ±22	914 ±15	+9	0.0695	0.71	1.326	2.9	0.1384	2.8	0.97
11.1	–	236	181	0.79	1.19	32	950 ±17	974 ±12	+3	0.0716	0.58	1.568	2.0	0.1588	1.9	0.96
12.1	0.01	752	5	0.01	0.92	68	648 ±9	660 ±9	+2	0.0616	0.43	0.899	1.6	0.1058	1.5	0.96
13.1	0.26	548	129	0.24	4.36	55	713 ±15	775 ±28	+8	0.0650	1.33	1.048	2.6	0.1170	2.2	0.86
14.1	0.06	197	115	0.60	0.45	27	965 ±17	983 ±15	+2	0.0719	0.72	1.601	2.1	0.1615	1.9	0.94
15.1	0.08	195	120	0.64	0.25	27	972 ±19	1009 ±15	+4	0.0728	0.73	1.634	2.2	0.1627	2.1	0.95
16.1	0.05	90	59	0.68	1.23	12	952 ±10	998 ±20	+5	0.0724	1.00	1.589	1.5	0.1591	1.2	0.76
17.1	0.41	215	85	0.41	1.63	21	682 ±7	961 ±24	+31	0.0711	1.20	1.095	1.6	0.1116	1.1	0.68
18.1	0.18	258	181	0.73	0.26	33	891 ±16	975 ±16	+9	0.0716	0.77	1.463	2.0	0.1482	1.9	0.93
22.1	0.04	869	8	0.01	0.69	78	640 ±8	660 ±10	+3	0.0616	0.44	0.887	1.4	0.1045	1.4	0.95
21.1	0.03	717	4	0.01	0.95	64	642 ±9	650 ±11	+1	0.0613	0.49	0.885	1.6	0.1048	1.5	0.95

Errors are 1-sigma; Pb_c and Pb^* indicate the common and radiogenic portions, respectively. Error in Standard calibration was 0.25% (not included in above errors but required when comparing data from different mounts) 0.1 – Common Pb corrected using measured ^{204}Pb . 2–Common Pb corrected by assuming $^{206}\text{Pb}/^{238}\text{U}$ – $^{207}\text{Pb}/^{235}\text{U}$ age-concordance.

Ma interval. Oliveira et al. (2015) interpreted the ages of 960–970 Ma as the maximum deposition age of the Marancó sedimentary protolith. These ages are similar to those described by Guimarães et al. (2012) in metavolcanic rocks intercalated in metasediments of the Riacho Gravatá Complex.

4.2.3. Riacho do Pontal domain

Two geochronological data are available for this domain (Fig. 1d). 1) U–Pb zircon LA-ICP-MS age of 1001 ± 4 Ma, obtained by Caxito et al. (2014) in a metric granite sill concordantly intruded within pelitic metarhytmite. 2) U–Pb zircon LA-ICP-MS age of 966 ± 5 Ma in the augen-gneisses of the Afeição suite (Caxito et al.,

2014), which are similar to those previously defined by Jardim de Sá (1994, 1988) using U–Pb zircon TIMS and Rb–Sr isochron age. The Afeição augen-gneisses are described (Caxito et al., 2012) as an NW–SE elongated body showing a conspicuous foliation, parallel to the main regional foliations. The central part of the intrusion is underformed and the analyzed sample was collected from this area. Caxito et al. (2012) suggest that the augen-gneisses of the Afeição suite are intrusive or thrust upon metasedimentary rocks, orthogneisses and migmatites of the basement and show tectonic contact with Brasileiro syn-collisional granitoids. These field relationships are similar to those described in the Cariris Velhos belt by Guimarães et al. (2012).




Fig. 5. a) Plot of $\text{Na}_2\text{O} + \text{K}_2\text{O} + \text{CaO}$ (MALI) versus SiO_2 (%wt) showing the distribution of the Tonian orthogneisses from the Borborema Province. Fields after [Frost and Frost \(2008\)](#) b) The composition range of the Tonian orthogneisses in the $\text{FeO}_{\text{tot}}/(\text{FeO}_{\text{tot}} + \text{MgO})$ vs. weight percent of SiO_2 diagram. Fields of ferroan and magnesian granitoids are from [Frost et al. \(2001\)](#).

4.2.4. Rio Preto domain

Some small plutons with Tonian ages occur in this domain. However, only one of them, the Serra da Pintada pluton ([Aquino and Batista, 2011](#)) has available geochronological data. Two other plutons have been described in regional map ([Aquino, 2014](#)), as of




Fig. 7. The Tonian orthogneisses in the ternary diagram $\text{Al}_2\text{O}_3/(\text{FeO} + \text{MgO}); 3^*\text{CaO}; 5^*(\text{K}_2\text{O}/\text{Na}_2\text{O})$, with fields representing composition of melts derived from a range of potential sources, after [Laurent et al. \(2014\)](#).

Tonian age, but geochronological data are not provided. The Serra da Pintada pluton comprises leucocratic, coarse-grained augen-gneisses similar to those of the Afeição suite in the Riacho do Pontal domain. Eleven zircon grains analyzed from one sample of the Serra da Pintada pluton ([Aquino and Batista, 2011](#)) gave a nice cluster of U–Pb zircon SHRIMP data, defining a concordia age of 969 ± 11 Ma ($\text{MSWD} = 0.72$), interpreted as the crystallization age of the orthogneiss protolith. Only two, of the thirteen analyzed zircon grains, have $^{208}\text{Pb}/^{238}\text{U}$ ages of 1.9 Ga and 2.1 Ga, suggesting some contribution of the Paleoproterozoic country rock orthogneisses in their source ([Aquino and Batista, 2011](#)).

5. Geochemistry

The Cariris Velhos granitoids described in this section include those from this work and those from the literature ie. the Cariris




Fig. 6. Shand's Index for the granitoids of the Tonian orthogneisses of the Borborema Province. Fields after [Maniar and Piccolli \(1989\)](#). CVB – Cariris Velhos belt; SI – Sítio do Icó; PEALD – Pernambuco Alagoas domain; LB – Lobo pluton; RC – Rocinha pluton; FL – Flores pluton; SD – Sergipano domain; SN – Serra Negra; PR – Poço Redondo migmatites; RPD = Riacho do Pontal domain.




Fig. 8. Chondrite normalized REE plots for the studied orthogneisses. Chondrite values from Nakamura (1974).

Velhos belt in the Transversal subprovince (Guimarães et al., 2012; Kozuch, 2003); from the South subprovince (Oliveira et al., 2010; Oliveira personal communication; Caxito et al., 2014; Cruz et al., 2014). However, there is no data available for the orthogneisses with crystallization ages younger than 900 Ma, and those from the Serra da Pintada pluton from the Rio Preto domain. The geochemical data from the Serra da Pintada augen - gneisses, we used the data presented in diagrams by Aquino and Batista (2011).

The Tonian augen-gneisses have a high SiO_2 (>71%). When migmatized the SiO_2 values vary widely but values lower <70 wt% dominate. The augen-gneisses plot mainly in the alkali-calcic and calc-alkaline fields of the MALI (Modified Alkali-Lime index) versus SiO_2 diagram of Frost and Frost (2008) (Fig. 5a). The migmatites plot in all fields of the MALI versus SiO_2 diagram as expected, due to the alkalis (K_2O and Na_2O) mobility during migmatization. They have high $\text{FeO}/(\text{FeO} + \text{MgO})$ ratios usually >0.8 and most of them are classified as Ferroan granitoids (Fig. 5b) in the $\text{FeO}/(\text{FeO} + \text{MgO})$ versus SiO_2 diagram (Frost et al., 2001). They range from slightly peraluminous to slightly metaluminous according to the Shand index (Fig. 6). According to the geochemical diagrams presented by Aquino and Batista (2011), the Serra da Pintada augen-gneisses have high SiO_2 , K_2O and low MgO contents. Their classification into magnesian or ferroan granitoids (Frost et al., 2001) is not possible, because Fe contents are not provided. However, from the diagrams, it is possible to know that the majority of them have FeO/MgO ratios >20, which point to ferroan granitoids.

In the ternary diagram $\text{Al}_2\text{O}_3/(\text{FeO} + \text{MgO}) - 3^*\text{CaO} - 5^*(\text{K}_2\text{O}/$

$\text{Na}_2\text{O})$, with fields representing compositions of potential sources (Laurent et al., 2014), the studied granitoids show an origin involving melting of distinct lithologies (Fig. 7). However, the Rocinha, Serra Negra, and Flores orthogneisses have composition point to an origin through melting of metasediments. The most part of the Recanto orthogneiss samples and few samples of the Poço Redondo migmatites fall in the field of metasedimentary source. All the other orthogneisses and migmatites plot in all fields of igneous source rocks, with composition of high-K mafic rocks (mainly Afeição augen-gneisses), low - K mafic rocks (few samples) and tonalities (most of the Cariris Velhos belt samples and Lobo orthogneisses). The Poço Redondo samples plot in the fields of high-Ca (mafic rocks) and low-Ca (metasedimentary) of the $\text{Al}_2\text{O}_3/(\text{FeO} + \text{MgO}) - 3^*\text{CaO} - 5^*(\text{K}_2\text{O}/\text{Na}_2\text{O})$ diagram. This geochemical behavior suggests that the Poço Redondo migmatites represent distinct rocks, migmatized during the Ediacaran period. The Poço Redondo sample with available geochronological data, plot into the field of magmas originated by melting of metasedimentary rocks, similar to those recorded in the Serra Negra augen - gneisses (Fig. 7).

Trace element variations within and between the studied plutons are likewise extensive. They show low Sr (20–150 ppm) and Nb (12–22 ppm) contents, variable Ba (100–1260 ppm), Rb (164–400 ppm) and Zr (144–408 ppm) contents and high abundances of Y (41–80 ppm) and Ga (19–30 ppm).

The chondrite normalized REE patterns for the studied granitoids from all domains show many similarities (Fig. 8a–d). They are




Fig. 9. The spidergram patterns of the studied Tonian orthogneisses, normalized to the values suggested by Thompson (1982), from the Transversal (a) and Southern subprovinces (b, c, d). In c) age of Poço Redondo from Carvalho (2005) and in d) ages from Caxito et al. (2014).

characterized by negative Eu anomalies and low $(Ce/Yb)_N$ ratios. The Lobo orthogneisses have the lowest, while the Flores orthogneisses have the highest total REE contents (Fig. 8b). The deeper negative Eu anomalies, with Eu/Eu^* ranging from 0.12 to 0.4, were recorded in the Pernambuco – Alagoas domain. The Serra da Pintada augen-gneisses have REE patterns (Fig. 8d) similar to those recorded in the orthogneisses from the Cariris Velhos belt and Afeição suite of the Riacho do Pontal domain.

The spidergram patterns of the orthogneisses from all domains normalized to the values suggested by Thompson (1982) (Fig. 9 a–d) show many similarities: troughs at Ba, Sr, P and Ti and, variable troughs at Nb and Ta. The Lobo orthogneisses also show troughs at Ba, Ti, Nb and Ta, however, they differ from the others orthogneisses patterns by troughs at Th and no troughs at Sr (Fig. 9b). They have patterns similar to those recorded in others fine-grained orthogneisses as Sitio do Icó, and some of the Poço Redondo migmatites. However, the dated sample from the Poço Redondo migmatite, has spidergram pattern similar to those recorded in the Serra Negra augen-gneisses (Fig. 9c). The spidergram patterns recorded in the studied augen-gneisses and most of the migmatites are similar to those recorded in extension-related granitoids (Thompson et al., 1984). The Serra da Pintada augen-

gneisses have primitive mantle spidergram patterns characterized by deep troughs at Sr, P and Ti, smaller troughs at Ba and peaks and smaller troughs at Nb (Aquino and Batista, 2011). These patterns are also similar to those recorded in extension-related granitoids.

The majority of the studied granitoids, including those from Serra da Pintada (see Aquino and Batista, 2011), plot in the A-type granites field (Fig. 10a) in the (FeO/MgO) versus $(Zr + Nb + Ce + Y)$ tectonic discriminant diagram of Whalen et al. (1987). However, some samples plot in the field of fractionated granites, while the fine-grained migmatite sample from Sitio do Icó, and another one dated at ~ 1.0 from Afeição suite, fall in the field of unfractionated M-, I- and S-type granites. The A-type Tonian granitoids are, with rare exception, distinct from OIB (Fig. 10b) in the Y/Ta versus Y/Nb diagram (Eby, 1992). This chemical signature is typical of A_2 -type granitoids (Eby, 1992). The A_2 -type signature is also recorded in the majority of the studied granitoids, when they are plotted in the $Nb-Y-Z/4$ and $Nb-Y-Ce$ diagrams (Fig. 10c,d) with fields after Eby (1992). However, although the Flores granitoids fall in the A_2 -type granites field in the $Nb-Y-Ce$ diagram, their high-Y and Nb contents are typical of A_1 -type granitoids.

The Sitio do Icó analyzed migmatite sample is magnesian, and plot within the calc-alkaline field of the MALI versus SiO_2 diagram




Fig. 10. The Tonian orthogneisses in the tectonic discriminant diagram of a) Whalen et al. (1987); and b) The Yb/Ta versus Yb/Nb diagram with fields of ocean island basalts (OIB) and island arc basalts (IAB) (Eby, 1992), and c) and d) A1 and A2 subgroups discrimination of A-type granites after Eby (1992). CVB – Cariris Velhos belt; SI – Sítio do Icó; PEALD – Pernambuco Alagoas domain; LB – Lobo pluton; RC – Rocinha pluton; FL – Flores pluton; SD – Sergipano domain; SN – Serra Negra; PR – Poço Redondo migmatites; RPD = Riacho do Pontal domain.

(Fig. 5a). It is distinct from the other Tonian granitoids from the Borborema Province, because it has low SiO_2 and HFSE contents. The analyzed sample shows spidergram pattern (Fig. 8a) with smaller troughs at Sr and Ti and lower HSFE contents, compared to the augen-gneisses. There is no geochemical data available for the Tonian orthogneisses from the east part of the PEAL domain and also for those of crystallization age younger than 0.9Ga.

6. Sm–Nd data

The Transversal subprovince concentrated most of the Sm–Nd whole rock data for the Tonian granitoids. The majority of them were obtained at the University of Kansas (Kozuch, 2003; Van Schmus et al., 2011), although a few were done at the Institute of Geosciences, São Paulo University (USP) and at the Geochronological Laboratories, Brasilia University.

The Tonian orthogneisses of the Cariris Velhos belt (Transversal subprovince) show Nd T_{DM} model ages ranging from 1.98 to 1.24Ga (Kozuch, 2003; Van Schmus et al., 1995, 2011). However, the majority of them have T_{DM} model ages within the 1.43–1.76 Ga interval (Fig. 11a,b). Those with T_{DM} model ages >1.8 Ga were recorded in the southwest part of the Cariris Velhos belt. Younger

T_{DM} model ages were recorded in the Sítio do Icó migmatized orthogneisses ($T_{\text{DM}} = 1.23\text{Ga}$ with $\epsilon\text{Nd}(t)$ of 2.89) and in the Serra Talhada orthogneisses (1.39 Ga and $\epsilon\text{Nd}(t)$ of 0.74).

In the Sergipano domain, Carvalho (2005) and Silva Filho et al. (1997) reported T_{DM} model ages ranging from 1.75 to 1.48Ga and $\epsilon\text{Nd}(t)$ values varying from 1.01 to –1.40 for the Poço Redondo migmatites. The Serra Negra pluton has T_{DM} model ages ranging from 1.53 to 1.66 Ga, and $\epsilon\text{Nd}(t)$ of –1.05 to –2.48 (Carvalho, 2005; Oliveira et al., 2010; Silva Filho et al., 1997).

Within the Pernambuco Alagoas domain, the Rocinha orthogneisses have T_{DM} model ages of 1.5Ga and $\epsilon\text{Nd}(t)$ value of –0.8 (Cruz and Accioly, 2013). Data from Silva Filho et al. (2014b) for migmatized orthogneisses (samples SI-100 and 125) show T_{DM} model ages ranging from 1.44 to 1.16Ga and ϵNd (980Ma) values ranging from 0.24 to 2.93.

The Sm–Nd data from the Riacho do Pontal domains, including migmatites and augen gneisses, show T_{DM} model ages ranging between 1.2 and 1.5 Ga, with $\epsilon\text{Nd}(t)$ from –1.0 to 3.1 (Caxito et al., 2014; Van Schmus et al., 1995, 2011).

Sm–Nd T_{DM} model ages recorded in the Tonian metaplutonic rocks (Fig. 11a,b) suggest participation of older crustal component in the genesis of their magmas. As pointed out by Van Schmus et al.




Fig. 11. Nd evolution paths for orthogneisses of the Tonian ages from the Borborema Province. a) CVB – Cariris Velhos belt data are from Van Schmus et al. (2011), Kozuch (2003). PEALD – Pernambuco Alagoas domain data are from (1) Cruz et al. (2014); (2) Silva Filho et al. (2014a,b). b) SD – Sergipano domain (Carvalho, 2005; Oliveira et al., 2010; Silva Filho et al., 1997). Depleted mantle curve of De Paolo (1981). Dashed line shows the upper limit of depleted mantle curves proposed by various authors (see Van Schmus et al., 2011).

(2011), large degrees of partial melting of a source having Sm/Nd model ages of 1.4–1.9 Ga (Mesoproterozoic lithosphere) can respond for such T_{DM} model ages recorded in the Tonian granitoids. However, this hypothesis is defeated by Van Schmus et al. (2011), using the argument that rocks with ages within this interval have a restrict occurrence in the Borborema Province and, that many of the plutons are hosted by Paleoproterozoic basement, with older Sm–Nd model ages (2.1–2.7 Ga). Another evidence that the Mesoproterozoic T_{DM} model ages of the Tonian granitoids reflect an old crustal component in their source, comes from the U–Pb data from the Riacho do Forno analyzed sample (Fig. 3). This sample has Paleoproterozoic zircon grains and Sm–Nd T_{DM} model age of 1.61Ga (Van Schmus et al., 2011). Thus, the Sm–Nd signatures recorded in the Tonian granitoids appear to reflect magmas formed by mixing of juvenile magmas within the 925 to 1000 Ma interval, with magmas generated by varying degrees of partial melting of Paleoproterozoic to Archean crust.

7. Discussions

The majority of the Tonian orthogneisses of the Borborema Province are ferroan granitoids. Ferroan magmas are formed by extreme differentiation or partial melting of tholeiitic to alkaline mafic magmas under reducing and relatively dry conditions in intraplate environments (Loiselle and Wones, 1979; Frost et al., 2001, 2008). The ferroan rocks in the SiO_2 versus MALI diagram are mostly alkali, although they can be alkali-calcic or even calc-alkaline as those from the Lachlan Fold Belt, described by Collins et al. (1982) and King et al. (2001). The ferroan granitoids from the Lachlan Fold Belt are aluminous (King et al., 1997) as the Tonian orthogneisses from the Borborema Province. Assimilation of high SiO_2 melts from the country rocks drives the magma to more calcic and peraluminous compositions (Frost and Frost, 2008).

The probably metasedimentary sources of the protoliths of the Serra Negra, Rocinha, Flores and some rocks of the Poço Redondo




Fig. 12. - Spidergram pattern of the Sitio do Icó sample compared to the patterns of Tonian metamafic volcanic rocks of the Riacho Gravata Complex (Guimarães et al., 2012) normalized to the MORB values (Pearce, 1983).

migmatites and Recanto orthogneisses are difficult to explain at present day. The metasediments in the Riacho Gravata Complex and Marancó subdomain have a Tonian maximum deposition age, and Tonian metamorphic event was not identified, up to now, in the Borborema Province. On the other hand, assimilation of deep-seated Paleoproterozoic sediments or melts produced by their fusion, could respond for the metasedimentary signature of these granitoids. The temperature required to produce an extractable magma may have been produced by either, mantle upwelling or mafic magmas influx during extension. The T_{DM} model ages >1.5 Ga, recorded in these granitoids, require some juvenile magma involved in their protolith.

Small degree of melting of such metasediments can produce high-SiO₂ melts, which were incorporated into juvenile Tonian magmas, increasing Ca- and Al-contents in the magma (Frost and Frost, 2008). Because, the studied granitoids are Ca-poor, plagioclase should be an important fractionated phase during the magma evolution, explaining in this way, the low Sr contents, and deep negative Eu anomalies recorded in these rocks, and explaining the Sm–Nd signature of these granitoids.

The Sitio do Icó migmatized orthogneisses sample is magnesian, and show mineralogy (high amount of modal amphibole) and geochemical signature distinct from others Tonian gneisses. Magnesian rocks are granitoids with composition ranging from calcic to alkali-calcic and are formed in arcs and post-collisional environments (Frost and Frost, 2008). However, magmas generated by partial melting of magnesian crustal rocks can inherit magnesian signature (Frost and Frost, 2008). The positive $\epsilon(Nd)$ values, T_{DM} model age of 1.23 Ga, and lack of rocks with ages ~ 1.2 Ga within the Borborema Province, do not support the hypothesis of a magma originated by melting of continental crustal rocks. It appears to have been originated from a depleted mantle source that underwent some crustal contamination. In Fig. 12, the similarities between the spidergram pattern normalized to the MORB values of this sample and the metavolcanic mafic rocks from the Riacho Gravata Complex of the Cariris Velhos belt, suggest that this rock is a metavolcanic migmatized rock.

According to the evolution model of Oliveira et al. (2010) for the

Sergipano domain, the Poço Redondo migmatites protoliths are arc-related, while the Serra Negra orthogneisses are A-type granites. However, the dated migmatite sample and the Serra Negra orthogneisses share similar extension-related geochemical signature (Fig. 9). On the other hand, the Tonian granitoids were intruded as small plutons and the Poço Redondo migmatites comprise a belt of more than 8 km wide and more than 30 km long. As pointed out before, the available geochemical data show that the analyzed samples represent rocks of distinct compositions and probably distinct sources. The lack, up to now, of evidence of a Tonian metamorphic event, associated to the geochemical similarities between the dated sample from the Poço Redondo migmatites and the Serra Negra pluton, do not support the hypothesis that the Poço Redondo migmatites are arc-related.

8. Conclusions

Tonian granitoids constitute the main magmatic expression of the Cariris Velhos belt of the Transversal subprovince and occur in minor volume in all domains of the South subprovince of the Borborema Province. However, they are absent, or not yet recognized, in the North subprovince.

The crystallization ages of the Tonian orthogneisses protoliths show that the main granitic pulse occurred in between 925 Ma to 970 Ma. The fine-grained migmatized rocks show older crystallization ages (~ 1.0 Ga), similar to those recorded in felsic metavolcanic rocks of the Riacho Gravata Complex. It suggests that most of the fine-grained migmatites can be metavolcanic felsic rocks. Thus, detailed work in these migmatites is necessary to better define their sources.

The studied granitoids have extension-related geochemical signature and most of them were generated by melting of igneous source rocks. However, the granitoids from the Sergipano domain and some from the south part of the Pernambuco – Alagoas domain appear to have a sedimentary component in their source. Interaction between juvenile magmas and melts generated by melting (or assimilation) of deep-seated Paleoproterozoic sediments, followed by plagioclase fractionation, is a plausible interpretation for the

genesis of the Tonian granitoids from the Sergipano and the south part of the PEAL domain. The Sm–Nd isotopic signatures also support this hypothesis. The heat sources is inferred to be mafic melts associated with crustal thinning. It is also important to emphasize the sediments input in the granitoids sources from south part of the Pernambuco – Alagoas domain and north part of the Sergipano domain.

However, the Cariris Velhos event is still open to significant debate. More detailed geochronological and geochemical data from the South subprovince, and detailed field works in the most migmatized area are necessary to understand the tectonic meaning of this event.

The Riacho do Pontal (Afeição suite) and Rio Preto (Serra da Pintada pluton) Tonian orthogneisses have geochemical signatures and crystallization ages similar to those recorded in the Cariris Velhos belt from the Transversal subprovince. These evidence associated to the dextral sense of the Pernambuco shear zone, suggest that during the Tonian period they were part of the same belt, which was disrupted by the dextral Pernambuco shear zone during the Brasiliano orogeny. Van Schmus et al. (2011) already proposed a link between the Transversal subprovince and the Riacho do Pontal domain.

Funding

IPG thanks CAPES for a post-doctoral scholarship (BEX 4127-0702). Grants from PRONEX/FACEPE program (APQ-0479-1.07/06), and from CNPq (477.462/2012-0) made possible IPG and AFSF to perform SHRIMP analyses at the ANU, Canberra, Australia.

Acknowledgments

GL thanks Dr Elton Dantas (Brasília University) for some U–Pb analytical data. The manuscript has been improved by the constructive review of the J. South Am. Earth Sci. reviewers, Dr. Evandro Klein (CPRM-Geological Survey of Brazil), Dr. Sanjeet K. Verma (Campinas University) and Dr. Lauro Nardi (Rio Grande do Sul Federal University).

References

- Accioly, A.C.A., Santos, C.A., Santos, E.J., Brito Neves, B.B., Rodrigues, J.B., McReath, I., 2010. Geochronology and geochemistry of the meta-volcanic rocks from riacho do tigre complex, borborema province - northeastern Brazil. In: VII SSAGI South American Symposium on Isotope Geology, Brasília (Abstract – CD-ROM).
- Accioly, A.C., 2003. A. Geologia, geoquímica e significado tectônico do Complexo Metanortosítico de Passira: Província Borborema, nordeste brasileiro. Doctoral thesis. São Paulo University, 168pp. (in Portuguese).
- Almeida, F.F.M., Hasui, Y., Brito Neves, B.B., Fuck, R.A., 1981. Brazilian structural provinces: an introduction. *Earth Sci. Rev.* 17, 1–21.
- Angelim, L.A.A., Kosin, M., 2001. Programa de Levantamentos Geológicos do Brasil. In: Folha Aracaju, N.W. (Ed.), Nota Explicativa (In Portuguese), CPRM, CDROM, Brasília, Brazil.
- Angelim, L.A.A., 1988. Programa Levantamentos Geológicos Básicos Do Brasil-plgb, Carta Geológica, Carta Metalogenética. Escala 1:100000 Folha SC.24-V-A-III, Santa Filomena, Estados de Pernambuco e Piauí. DNP/CPRM, 146 pp.
- Aquino, M.R., Batista, I.H., 2011. In: Geologia e recursos minerais da folha Avelino Lopes – SC.23-Z-A-III. Programa de Levantamentos Geológicos do Brasil. CPRM online. www.cprm.gov.br. Teresina, Brasil. Nota Explicativa. 88pp. (in Portuguese).
- Aquino, M.R., 2014. Carta Geológica – Folha Curimatá, Escala 1:100.000 Programa Geologia Do Brasil. online. www.cprm.gov.br (legend in Portuguese).
- Bittar, S.M.B., 1998. Faixa Pianco – Alto Brigida: Terrenos tectono estratigraficos sobregimes metamórficos e deformacionais contrastantes, regimes metamórficos e deformacionais contrastantes. Doctoral thesis. Instituto de Geociências, Universidade de São Paulo, 126pp. (in Portuguese).
- Brito, M.F.L., Freitas, S., 2011. Caracterização petrológica e geotectônica do ortognaisse Lobo no domínio Pernambuco-Alagoas, Província Borborema: Nordeste do Brasil. In: XIII Brazilian Geochemical Congress. Gramado-rs, Brazil. Extended Abstract – CD-Rom, pp. 796–799 (in Portuguese).
- Brito, M.F.L., Mendes, V.A., Paiva, I.P., 2008. Metagranitóide Serra das Flores: magmatismo Toniano (tipo-A) no Domínio Pernambuco-Alagoas, Nordeste do Brasil. In: 44° Brazilian Geological Congress, Curitiba. Abstract in CD-ROM (in Portuguese).
- Brito Neves, B.B., Campos Neto, M.C., Van Schmus, W.R., Santos, E.J., 2001. O Sistema Pajeu–Paraíba e o Maciço São Jose de Campestre no leste da Borborema. *Rev. Bras. Geoc.* 31, 173–184 (in Portuguese).
- Brito Neves, B.B., Santos, E.J., Schmus, W.R.V., 2000. Tectonic history of the Borborema Province. In: Cordani, Umberto, Milani, Edson Jose, Thomaz Filho, Antonio, Campos, Diogenes de Almeida (Eds.), Tectonic Evolution of South America. Rio de Janeiro: 31st International Geological Congress, pp. 151–182. Special Publication.
- Brito Neves, B.B., Cordani, U.G., 1991. Tectonic evolution of South America during the late Proterozoic. *Precambrian Res.* 53, 23–40.
- Brito Neves, B.B., 1983. O Mapa Geológico Do Nordeste Oriental Do Brasil, 1:1,000,000. Thesis of Livre Docência. São Paulo University, 177pp. (in Portuguese).
- Carvalho, M.J., 2005. Tectonic Evolution of the Marancó-poço Redondo Domain: Records of the Cariris Velhos and Brasiliano Orogenesis in the Sergipano Belt, NE Brazil. Doctoral thesis. Campinas University, 202pp. (in Portuguese).
- Caxito, F.A., Uhlein, A., Sanglard, J.C.D., Gonçalves-Dias, T., Mendes, M.C.O., 2012. Depositional systems and stratigraphic review proposal of the Rio Preto Fold Belt, northwestern Bahia/southern Piauí. *Rev. Bras. Geoc.* 42 (3), 523–538.
- Caxito, F.A., Uhlein, A., Dantas, E.L., 2014. The Afeição augen-gneiss Suite and the record of the Cariris Velhos Orogeny (1000–960 Ma) within the Riacho do Pontal fold belt, NE Brazil. *Journal of South American Earth Sciences* 51, 12–27.
- Collins, W.J., Beams, S.D., White, A.J.R., Chappell, B.W., 1982. Nature and origin of A-type granites with particular reference to southeastern Australia. *Contrib. Min. Pet.* 80, 189–200.
- Cruz, R.F., Pimentel, M.M., Accioly, A.C.A., Rodrigues, J.B., 2014. Geological and isotopic characteristics of granites from the Western Pernambuco-Alagoas Domain: implications for the crustal evolution of the Neoproterozoic Borborema Province. *Brazilian Journal of Geology* 44 (4), 627–652.
- Cruz, R.F., Accioly, A.C.A., 2013. Petrografia, geoquímica e idade U/Pb do Ortognaisse Rocinha, no Domínio Pernambuco-Alagoas W da Província Borborema. *Estudos Geológicos* 23 (2), 3–27. www.ufpe.br/estudosgeologicos (in portuguese).
- Dantas, E.L., 1997. Geocronologia U-Pb e Sm-Nd de terrenos arqueanos e paleoproterozoicos do Maciço Caldas Brandão, NE do Brasil. Doctoral thesis. Paulista State University (UNESP), 206pp. (in Portuguese).
- Davison, I., Souza, R., 1989. Tectonic evolution of the Sergipano fold Belt, NE Brazil, during the Brasiliano orogeny. *Precambrian Res.* 45, 319–342.
- Dias, R.C., Juliá, J., Schimmel, M., 2015. Rayleigh-wave, group – velocity tomography of the Borborema Province, NE Brazil, from ambient seismic noise. *Pure Appl. Geophys.* 172, 1429–1449.
- Eby, G.N., 1992. Chemical subdivision of the A-type granitoids: petrogenetic and tectonic implications. *Geology* 20, 641–644.
- Egydio-Silva, M., Karmann, I., Trompette, R.R., 1989. Litoestratigrafia do Supergrupo Espinhaço e Grupo Bambuí no noroeste do estado da Bahia. *Rev. Bras. Geoc.* 19 (2), 101–112 (in Portuguese).
- Fetter, A.H., 1999. U-pb and Sm-nd Constraints on the Crustal Framework and Geologic History of Ceará State, NW Borborema Province, NE Brazil: Implications for the Assembly of Gondwana. PhD thesis. Kansas University 164pp.
- Frost, B.R., Frost, C.D., 2008. A geochemical classification for Feldspathic igneous rocks. *J. Pet.* 49, 1955–1969.
- Frost, B.R., Barnes, C., Collins, W., Arculus, R., Ellis, D., Frost, C., 2001. A chemical classification for granitic rocks. *J. Pet.* 42, 2033–2048.
- Gomes, F.E.M., Vasconcelos, A.M., 1991. Programa Levantamento Geológicos Básicos Do Brasil. Carta Geológica, Carta Metalogenética, Escala 1:100000. Folha SC.24-V-A-II, Paulistana, Estados de Pernambuco e Piauí. DNP/CPRM, 146 pp.
- Guimarães, I.P., Van Schmus, W.R., Brito Neves, B.B., Bittar, S.M., Silva Filho, A.F., Armstrong, R., 2012. U Pb zircon ages of orthogneisses and supracrustal rocks of the Cariris Velhos belt: onset of Neoproterozoic rifting in the Borborema Province, NE Brazil. *Precambrian Res.* 192–195, 52–77.
- Jardim de Sá, E.F., 1994. A Faixa Seridó (Província Borborema, NE do Brasil) e o seu significado geodinâmico na cadeia Brasileira/Pan Africana. Doctoral thesis. Brasília University 803pp. (in Portuguese).
- Jardim de Sá, E.F., Macedo, M.H.F., Fuck, R.A., Kawashita, K., 1992. Terrenos proterozoicos na província Borborema e a margem norte do Cráton do São Francisco. *Rev. Bras. Geoc.* 22, 472–480.
- Jardim de Sá, E.F., Macedo, M.H.F., Torres, H.H.F., Kawashita, K., 1988. Geochronology of metaplutonics and the evolution of supracrustal belts in the Borborema province. In: VII Latino-americano Geological Congress. Extended Abstract, pp. 49–62.
- King, P.L., Chappell, B.W., Allen, C.M., White, A.J.R., 2001. Are A-type granites the high-temperature felsic granites? Evidence from fractionated granites of the Wangrah Suite. *Aust. J. Earth Sci.* 48, 501–514.
- King, P.L., White, A.J.R., Chapell, B.W., Allen, C.M., 1997. Characterization and origin of aluminous A-type granites from the Lachlan Fold Belt, southeastern Australia. *J. Pet.* 38, 371–391.
- Kozuch, M., 2003. Isotopic and Trace Element Geochemistry of Early Neoproterozoic Gneiss and Metavolcanic Rocks in the Cariris Velhos Orogen of the Borborema, Gneissic and Metavolcanic Rocks in the Cariris Velhos Orogen of the Borborema Province, Brazil and Their Bearing Tectonic Setting. PhD thesis. Kansas University, 199pp.
- Lages, G.A., Dantas, E.L., Santos, L.C.M., 2014. Sítio do icó orthogneiss, east of transversal zone, borborema province: stonian orthogneiss with juvenile characteristics. In: 9th South American Symposium on Isotope Geology.

- Abstract, 113.
- Laurent, O., Martina, H., Moyena, J.F., Doucelance, R., 2014. The diversity and evolution of late-Archean granitoids: Evidence for the onset of “modern-style” plate tectonics between 3.0 and 2.5 Ga. *Lithos* 205, 208–235.
- Leite, P.R.L.B., Bertrand, J.M., Lima, E.S., Leterrier, J., 2000. Timing of granitic magmatism in the northern Borborema Province, Brazil: a U-Pb study of granitoids from the Alto Pajeu Terrain. *J. South Am. Earth Sci.* 13, 549–559.
- Loiselle, M.C., Wones, D.R., 1979. Characteristics and origin of anorogenic granites. *Geol. Soc. Amer. Abst. Prog.* 11, 468.
- Maniar, P.D., Piccoli, P.M., 1989. Tectonic discrimination of granitoids. *Geol. Soc. Am. Bull.* 101, 635–643.
- Moraes, J.F.S., 1992. Petrologia das rochas máficas-ultramáficas da seqüência vulcanosedimentar de Monte Orebe, PE-PI. M.S.dissertation. Bahia Federal University, 98 pp. (in Portuguese).
- Nakamura, N., 1974. Determination of REE, Ba, Fe, Mg, Na and K in carbonaceous and ordinary chondrite. *Geochim. Cosmochim. Acta* 38, 757–775.
- Neves, S.P., 2003. Proterozoic history of the Borborema province (NE Brazil): correlations with neighboring Cratons and Pan-African belts and implications for the evolution of western Gondwana. *Tectonics* 22, 1031–1044.
- Neves, S.P., Bruguier, O., Vauchez, A., Bosch, D., Silva, J.M.R., Mariano, G., 2006. Timing of crust formation, deposition of supracrustal sequences, and Transamazonian and Brasiliano metamorphism in the East Pernambuco belt (Borborema Province, NE Brazil): Implications for western Gondwana assembly. *Precambrian Res.* 149, 197–216.
- Neves, S.P., Monie, P., Bruguier, O., da Silva, J.M.R., 2012. Geochronological, thermochronological and thermobarometric constraints on deformation, magmatism and thermal regimes in eastern Borborema Province (NE Brazil). *J. South Am. Earth Sci.* 38, 129–146.
- Neves, S.P., Lages, G.A., Brasilino, R.G., Miranda, A.W.A., 2015. Paleoproterozoic accretionary and collisional processes and the build-up of the Borborema Province (NE Brazil): geochronological and geochemical evidence from the Central Domain. *J. South Am. Earth Sci.* 58, 165–187.
- Oliveira, R.G., 1998. Arcabouço geotectônico da região da Faixa Riacho do Pontal, Nordeste do Brasil: dados aeromagnéticos e gravimétricos. Masters Dissertation. São Paulo University, 157pp. (in Portuguese).
- Oliveira, E.P., Windley, B., Araujo, M.N.C., 2010. The Neoproterozoic Sergipano orogenic belt, NE Brazil: a complete plate tectonic cycle in western Gondwana. *Precambrian Res.* 181, 64–84.
- Oliveira, E.P., Bueno, J.F., McNaughton, N.J., Silva Filho, A.F., Nascimento, R.S., Donatti-Filho, J.P., 2015. Age, composition, and source of continental arc- and syn-collision granites of the Neoproterozoic Sergipano Belt, Southern Borborema Province, Brazil. *J. South Am. Earth Sci.* 58, 258–280.
- Paces, J.B., Miller Jr., J.D., 1993. Precise U–Pb ages of Duluth complex and related mafic intrusions, mafic intrusions, northeastern Minnesota: geochronological insights into physical, petrographic paleomagnetic, and tectonomagmatic processes associated with the 1.1 Ga midcontinental rift system. *J. Geophys. Res.* 98, 13997–14013.
- Pimentel, M.M., Fuck, R.A., 1992. Neoproterozoic crustal accretion in central Brazil. *Geology* 20, 375–379.
- Pearce, J.A., 1983. Role of the sub-continental lithosphere in magma genesis at active continental margins. In: Hawkesworth, C.J., Norry, M.J. (Eds.), *Continental basalts and Mantle Xenoliths*. Shiva Geology Series, 230–249.
- Pimentel, M.M., Whitehouse, M.J., Viana, M.G., Fuck, R.A., Machado, N., 1997. The Mara Rosa arc in the Tocantins province: further evidence for Neoproterozoic crustal accretion in central Brazil. *Precambrian Res.* 81, 299–310.
- Pimentel, M.M., Fuck, F.A., Jost, H., Ferreira Filho, C.F., Araújo, S.M., 2000. The basement of the Brasília fold belt and the Goiás magmatic arc. In: Cordani, U., Milani, E.J., Thomaz Filho, A., Campos, D.A. (Eds.), *Tectonic Evolution of South America*. 31st International Geological Congress, Rio de Janeiro, Brazil, pp. 195–229.
- Rogers, J.J.W., Santosh, M., 2002. Configuration of Columbia, a Mesoproterozoic supercontinent. *Gondwana Res.* 5, 5–22.
- Rogers, J.J.W., Santosh, M., 2003. Supercontinents in Earth history. *Gondwana Res.* 6, 357–368.
- Rogers, J.J.W., Santosh, M., 2004. *Continents and Supercontinents*. Oxford University Press, Oxford.
- Sá, J.M., Bertrand, J.M., Leterrier, J., Macedo, M.H.F., 2002. Geochemistry and geochronology of pre-Brasiliano rocks from the transversal zone, borborema province, northeast Brazil. *J. South Am. Earth Sci.* 14, 851–866.
- Santos, E.J., 1995. O complexo granítico Lagoa das Pedras: Acreção e colisão na região de Floresta (Pernambuco) Província da Borborema. Doctoral thesis. São Paulo University, 220pp.
- Santos, E.J., Van Schmus, W.R., Brito Neves, B.B., Oliveira, R.G., Medeiros, V.C., 1999. Terranes and their boundaries in the Proterozoic borborema province, northeast Brazil. In: I International Symposium on Tectonics of the Brazilian Geologic Society, and VII SNET, Bahia, Brazil, Extended Abstracts, pp. 120–124.
- Santos, E.J., Van Schmus, W.R., Kozuch, M., Brito Neves, B.B., 2010. The cariris velhos tectonic event in northeast Brazil. *J. South Am. Earth Sci.* 29, 61–76.
- Sato, K., Basei, M.A.S., Siga, O.J., 2008. Novas técnicas aplicadas ao método U-Pb no CPGeo - IGC/USP: avanços na digestão química, espectrometria de massa (TIMS) e exemplos de aplicação integrada com SHRIMP [New techniques applied to U-Pb method at CPGeo - Igc/USP: advances in chemical digestion, mass spectrometry (TIMS) and examples of application integrated with SHRIMP]. *Geol. Usp. Série Científica* 8, 77–99 (in Portuguese).
- Silva Filho, A.F., Guimarães, I.P., Van Schmus, W.R., Armstrong, R.A., Silva, J.M.R., Osaco, L.S., Concentino, L.M., 2014a. SHRIMP U–Pb zircon geochronology and Nd signatures of supracrustal sequences and orthogneisses constrain the Neoproterozoic evolution of the Pernambuco–Alagoas domain, southern part of Borborema Province, NE Brazil. *Inter. J. Earth Sci.* 103 (8), 2155–2190.
- Silva Filho, A.F., Guimarães, I.P., Van Schmus, W.R., Armstrong, R.A., Santos, L.S., Concentino, L.M., Lima, D., 2014b. Shrimp oxygen, U-Pb, and Hf data of Tonian orthogneisses, evidence of juvenile crust in the Pernambuco – Alagoas domain of the Borborema Province, NE Brazil. In: 9th South American Symposium on Isotope Geology. Program and Abstracts, 95.
- Silva Filho, A.F., Guimarães, I.P., Brito, M.F.L., Pimentel, M.M., 1997. Geochemical signatures of main Neoproterozoic late-tectonic granitoids from the Sergipano Fold Belt, Brazil: Significance for the Brasiliano Orogeny. *Inter. Geol. Rev.* 39 (7), 639–659.
- Souza, Z.S., Martin, H., Peucat, J.-J., Sá, E.F.J., Macedo, M.H.F., 2007. Calc-alkaline magmatism at the archaean – Proterozoic transition: the Caico complex basement (NE Brazil). *J. Pet.* 48, 2149–2185.
- Stern, R.A., 1997. The GSC sensitive high resolution ion microprobe (SHRIMP): analytical techniques of zircon U–Th–Pb age determinations and performance evaluation. In: *Radiogenic age and isotopic studies: Report 10*. *Geol. Surv. Can. Curr. Res.* 1–31.
- Thompson, R.N., 1982. Magmatism of the British tertiary volcanic province. *Scottish J. Geol.* 18, 50–107.
- Thompson, R.N., Morrison, M.A., Hendry, G.L., Parry, S.J., 1984. An assessment of the relative roles of crust and mantle in magma genesis: an elemental approach. *Phil. Trans. R. Soc. A310*, 549–590.
- Trompette, R., 1997. Neoproterozoic (~600 Ma) aggregation of Western Gondwana: a tentative scenario. *Precambrian Res.* 82, 101–112.
- Trompette, R., 2000. Gondwana evolution; its assembly at around ≈600 Ma. *Comptes Rendus Acad. Sci. Paris* 330, 305–315.
- Van Schmus, W.R., Brito Neves, B.B., Harckspacher, P., Babinsky, M., 1995. U/Pb and Sm/Nd geochronologic studies of the eastern Borborema Province, Northeastern Brazil: initial conclusions. *J. South Am. Earth Sci.* 8, 267–288.
- Van Schmus, W.R., Oliveira, E.P., Silva Filho, A.F., Toteu, S.F., Penaye, J., Guimarães, I.P., 2008. Proterozoic links between the borborema province, NE Brazil, and the Central african fold belt. *Geol. Soc. Lond.* 294, 69–99. Special Publication.
- Van Schmus, W.R., Kozuch, M., Brito Neves, B.B., 2011. Precambrian history of the Zona Transversal of the Borborema Province, NE Brazil: insights from Sm/Nd and U/Pb geochronology. *J. South Am. Earth Sci.* 31, 227–252.
- Whalen, J.B., Currie, K.L., Chappell, B.W., 1987. A-types granites: geochemical characteristics, discrimination and petrogenesis. *Contrib. Mineral. Pet.* 95, 407–419.
- Williams, I.S., 1998. U–Th–Pb geochronology by ion microprobe. In: McKibben, M.A., Shanks III, W.C., Ridley, W.I. (Eds.), *Applications of Microanalytical Techniques to Understanding Mineralizing Processes*. *Rev. Econ. Geol.* 7, pp. 1–35.