



Short communication

Bothrops erythromelas (Amaral, 1923) venom induces apoptosis on renal tubular epithelial cells



Fabíola Carine Monteiro de Sousa^a, Antônio Rafael Coelho Jorge^a,
Ramon Róseo Paula Pessoa Bezerra de Menezes^a, Alba Fabíola Costa Torres^b,
Clarissa Perdigão Mello^b, Danya Bandeira Lima^b, Diva Maria Borges Nojosa^c,
Alexandre Havt^a, Renata Sousa Alves^b, Alice Maria Costa Martins^{b,*},
Helena Serra Azul Monteiro^a

^a Department of Physiology and Pharmacology, Federal University of Ceara, Fortaleza, Ceara, Brazil

^b Department of Clinical and Toxicological Analysis, Federal University of Ceara, Fortaleza, Ceara, Brazil

^c Regional Nucleus of Ophiology (NUROF), Federal University of Ceara, Fortaleza, Ceara, Brazil

ARTICLE INFO

Article history:

Received 5 February 2016

Received in revised form

28 March 2016

Accepted 19 April 2016

Available online 23 April 2016

Keywords:

Bothropoides erythromelas venom

MDCK cells

Apoptosis

ABSTRACT

Bothrops erythromelas is responsible for a large number of snakebite incidents in Northeastern Brazil. Previously, we showed the effects of whole *B. erythromelas* venom in an isolated kidney model. To continue the study with *B. erythromelas* venom, the present work aims to study the effects of this venom on MDCK tubular epithelial cells and assess gene expression involved in kidney injury, aiming at elucidating the mechanisms responsible for renal toxicity. Cytotoxicity in MDCK cells showed an IC₅₀ of 93 µg/mL and predominant apoptotic involvement demonstrated by flow cytometry assays and expression of caspase-3 and caspase-8. In conclusion, we suggest that *Bothropoides erythromelas* venom causes apoptosis with involvement of the caspases, probably through the extrinsic pathway.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Ophidic accidents are an important public health problem in Brazil, with high morbidity and mortality (Warrell, 2004; Albuquerque et al., 2013). The genus *Bothrops* and *Bothropoides* are responsible for 70% of ophidic accidents in Brazil (Oliveira et al., 2010). Venom from species of *Bothrops* genus show variable composition and biological effects (Queiroz et al., 2008).

Bothrops erythromelas (Amaral, 1923) commonly known in Brazil as “Jararaca-da-seca” is responsible for a great deal of snakebites in Northeastern Brazil (Rocha, 2008). The envenomation causes hemorrhage, edema, pain, myonecrosis, hypotension and induces acute renal failure (Sanchez et al., 1992; Sgrignolli et al., 2011).

Acute kidney injury is one of the main causes of death in snakebite cases (Sgrignolli et al., 2011; Albuquerque et al., 2013; Martines et al., 2014) and its pathogenesis is not well understood. Previously, we showed the effects of whole *B. erythromelas* venom

in an isolated kidney model (Martins et al., 2005). To continue the study with *B. erythromelas* venom (BeV), this short communication aims to study the effects of this venom on MDCK distal tubular epithelial renal cells and assess gene expression involved in kidney injury, aiming at elucidating the mechanisms responsible for renal toxicity.

2. Materials and methods

2.1. Cell culture, venom and chemical compounds

Madin–Darby Canine Kidney (MDCK) epithelial cells were cultivated in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS) (Sigma) at 37 °C and 5% CO₂. BeV was collected from adult specimens maintained in the Regional Nucleus of Ophiology of Federal University of Ceara (NUROF-UFC), lyophilized and maintained at –20 °C until further use. RPMI medium and (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay were obtained from Sigma Chemical Co. (St. Louis, MO, USA), AnnexinV/FITC Apoptosis Detection Kit was obtained from BD Pharmingen (CA, USA).

* Corresponding author.

E-mail address: martinsalice@gmail.com (A.M.C. Martins).

2.2. Cytotoxicity assay

Mitochondrial functionality was measured by MTT colorimetric assay. MDCK cells were cultured in sterile 96-well microtiter plates and were left to adhere to the plate overnight. After that, they were treated with different concentrations of *Bothrops erythromelas* venom (100; 50; 25 and 12.5 $\mu\text{g}/\text{mL}$) and incubated at 37 °C for 24 h. MTT reagent (Sigma-Aldrich, St. Louis, MO, USA; 5 mg/mL in PBS) was added to each well and incubated for 4 h at 37 °C. Finally, the medium was removed and the precipitated formazan crystals were dissolved in 10% Sodium dodecyl sulphate (Vetec, Rio de Janeiro-RJ, Brazil) in HCl 0.01 N. After 17 h, absorbance at 570 nm was performed in a microplate reader (Biochrom® Asys Expert Plus). Cell viability was calculated in comparison with the control group. The IC₅₀ (venom concentration capable of inhibiting 50% of cell growth) was determined by non-linear regression (Mosmann, 1983).

2.3. Annexin V-FITC and 7AAD staining

Cells treated with IC₅₀ of BeV (93.3 $\mu\text{g}/\text{mL}$) incubated during 24 h were stained with fluorescein isothiocyanate (FITC)-conjugated to annexin V/7AAD according to the manufacturer's instructions (BD Pharmingen, CA, USA). The population of annexin V-7AAD viable cells was evaluated by flow-cytometry. Data were collected in a FACSCalibur (Becton-Dickinson, Mountain View, Calif.) and analyzed using Cell Quest software (Becton-Dickinson).

2.4. Gene expression pro-apoptotics and anti-apoptotic

The expression of pro-apoptotic (Caspase-3 and Caspase-8) and anti-apoptotic (*Mcl-1* and *Bcl-XL*) genes were analyzed using iQ5 Real Time PCR Detection System (Bio-Rad). The mRNA sequence of pro-apoptotic and anti-apoptotic genes was obtained from the website of the National Center for Biotechnology Information (NCBI). Primer design was performed using the Oligo Designer Perfect™ available on the Invitrogen® web site (<http://www.invitrogen.com>). To assess gene expression, MDCK cells were grown in 24-well culture plates at a concentration of 1×10^5 cells/mL. After 24 h of cultivation in an incubator with 5% CO₂ at 37 °C, BeV was added to each well at concentrations of 46.65 and 23.32 $\mu\text{g}/\text{mL}$ (respectively 1/2 and 1/4 of IC₅₀ of venom) in hexaplicate and the cells were incubated with 5% CO₂ at 37 °C. Doxorubicin (3.12 $\mu\text{g}/\text{mL}$) was used as a positive control. After 24 h, the cells were removed from the wells through exposure to trypsin-EDTA (0.25/0.02% v/v) and centrifuged. Six precipitates was obtained for each concentration of the test substance, which were mixed to obtain a single sample of each concentration. The following experiment was the isolation of total RNA using the RNeasy Mini kit (Qiagen) and Qiacube automation equipment.

2.5. Statistical analysis

All results were expressed as mean \pm standard error of mean (SEM). The results of the experiments ($n = 3$) were submitted to analysis of variance (ANOVA) and Bonferroni post-test with a level of significance of * $p < 0.05$.

3. Results and discussion

Previous studies showed that cell death plays an important role in the nephrotoxicity caused by snake venom from the *Bothrops* and *Bothropoides* genus (Morais et al., 2013; Mello et al., 2014; Marinho et al., 2015).

Previously, we demonstrated the effects of whole BeV on an

isolated kidney model and it was be able to decreased percentage tubular transport of sodium and potassium (Martins et al., 2005). Tubular epithelial cells are the main targets of acute injury kidney caused by venoms (Sitprija and Sitprija, 2012). The loss of tubular epithelial cell function induced by BeV could cause decrease on percentage tubular transport of electrolytes. To continue these studies, we investigated the effects of BeV on the renal tubular epithelial cells.

MDCK cells constitute a very-well established cell line and have been extensively employed in the investigation of several cell processes, including epithelial transport and cell response to toxic agents and venoms (Chan et al., 1989; Collares-Buzato et al., 1994, 1998; 2002; Schwerdt et al., 2004; Peixoto and Collares-Buzato, 2005; Chen et al., 2006; Nascimento et al., 2007; Kusma et al., 2008).

In this study, we evaluated the cytotoxicity of whole venom on epithelial distal tubular MDCK renal cells using MTT assay and observed an IC₅₀ = 93 $\mu\text{g}/\text{mL}$ (Fig. 1A). The results indicated that the venom of *B. erythromelas* is cytotoxic to the renal cells, being similar the results of Collares-Buzato et al. (2002) who investigated the *in vitro* effects of whole venom of *Bothrops moojeni* using MDCK cell culture.

Apoptotic and necrotic cells can be detected by analyzing their light-scattering properties in flow cytometry (Krysko et al., 2011). Annexin-V and 7AAD were used for cell staining to identify the cell death pathway. In MDCK cells, flow cytometric analysis of BeV showed that the apoptotic cell and late apoptotic cell populations are significantly increased after 24 h-exposure to the IC₅₀, with a small percentage of necrotic cells (Fig. 1B). Taken together, these results show that apoptosis is predominant.

The major compound of BeV is metalloproteinases (Jorge et al., 2015), a class of apoptosis-inducing snake venom protein, such as described by Brenes et al. (2010), Masuda et al. (1998, 2000, 2001). Furthermore, the *Bothropoides pauloensis* venom, of which main compound is metalloproteinases (Rodrigues et al., 2012) was also described as an inducer of apoptosis (Marinho et al., 2015).




Fig. 1. Cytotoxic effect of *Bothropoides erythromelas* venom on MDCK cells. (A) MDCK cells were treated with different concentrations of *B. erythromelas* venom for 24 h and were evaluated by the reduction in the 3-(4,5-Dimethylthiazol-2-Yl)-2,5-Diphenyltetrazolium Bromide (MTT) salt in cells under BeV concentration-curve after 24 h of treatment. **(B)** Cell death was measured by annexin V and 7AAD staining and detected by flow cytometry. MDCK cells were treated with IC₅₀ (93.31 $\mu\text{g}/\text{mL}$) of BeV after 24 h. All data are expressed as mean \pm SEM of three independent experiments with three replicates and compared with negative control (One-way analysis of variance and Bonferroni post-test, * $p < 0.05$).




Fig. 2. Expression of pro-apoptotic and anti-apoptotic genes. Expression of (A) Caspase-3, (B) Caspase-8, (C) *Bcl-XL* and (D) *Mcl-1* gene. MDCK cells were treated with BeV at concentrations of 46.65 µg/mL (1/2 IC₅₀) and 23.32 µg/mL (1/4 of IC₅₀) of venom in hexaplicate and after 24 h the following experiment was performed. Doxorubicin (3.12 µg/mL) was used as a positive control. All data are expressed as mean ± SEM of three independent experiments and compared with negative control (One-way analysis of variance and Bonferroni post-test, *p < 0.05).

A number of cytotoxic agents have been reported to induce apoptosis depending on the intensity of their deleterious effects on cells. The predominant characteristics of either one or the other type of cell death are usually determined by the intensity and not by the specificity of stimulus (Orrenius et al., 1992; Bonfoco et al., 1995). In general, apoptosis occurs after exposure to low doses of the toxic agent, while necrosis requires high doses (Morais et al., 2013; Mello et al., 2014).

In this study, we also evaluated the expression of pro-apoptotic and anti-apoptotic genes to assess the involved cell death. Doxorubicin (Doxo), used as control in this assays, induces apoptosis in several cell lines (Kim et al., 2009; Boccellino et al., 2010; Fan et al., 2010; Martirosyan et al., 2010; Wang et al., 2010).

The BeV stimulated the expression of caspase-3 at a concentration of 1/4 IC₅₀ and of caspase-8 at both concentrations (1/4 and 1/2 IC₅₀). These effects were similar to those exhibited by Doxorubicin (Fig. 2A and B). Caspase-8 initiates apoptosis by activating the effector caspase (caspase-3), which, once activated, cleaves key proteins that process the apoptosis. Caspase-8 activates the extrinsic pathway of apoptosis, while caspase-3 is common to both pathways (Parrish et al., 2013).

In this study, the expression of *Mcl-1* increased at concentration 46.65 µg/mL, when compared with the negative control (Fig. 2C), suppressing apoptosis only at the highest concentration. The whole venom did not alter the expression of *Bcl-XL* at the tested concentrations, when compared with the negative control (Fig. 2D). The mitochondrial pathway is often activated in response to DNA damage, involving the activation of a member of the pro-apoptotic *Bcl-2* family (Shamas-Din et al., 2013). The anti-apoptotic members of the *Bcl-2* family inhibit death by apoptosis by preventing the formation of pores in the mitochondrial membrane, thus inhibiting

the leakage of cytochrome c into the cytosol (Gottlieb et al., 2000; Landeta et al., 2011). Subsequent caspases are activated, culminating in the cleavage of specific substrates and cell death by apoptosis (Anuradha et al., 2001).

Apoptosis can be initiated in two basic ways, known as intrinsic or mitochondrial and extrinsic or through death receptors (Schmitz et al., 2000; Ferri and Kroemer, 2001a, 2001b; Baetu and Hiscott, 2002). According to current evidence, it has been proposed that apoptosis caused by BeV probably acts through the extrinsic pathway or through death receptors.

In conclusion, we suggest that *Bothrops erythromelas* venom causes apoptosis with involvement of the caspases, probably through the extrinsic pathway.

Acknowledgments

This project was supported by grants from Coordenação de Aperfeiçoamento do Ensino Superior (CAPES), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico do Estado do Ceará (FUNCAP).

Transparency document

Transparency document related to this article can be found online at <http://dx.doi.org/10.1016/j.toxicon.2016.04.040>.

References

- Albuquerque, P.L., Jacinto, C.N., Silva Junior, G.B., Lima, J.B., Veras Mdo, S., Daher, E.F., 2013. Acute kidney injury caused by *Crotalus* and *Bothrops* snake venom: a review of epidemiology, clinical manifestations and treatment. *Rev. Inst. Med.*

- Trop. Sao Paulo 55, 295–301.
- Amaral, A., 1923. New genera and species of snakes. *Proc. N. Engl. Zool. Club* 8, 85–105.
- Anuradha, C.D., Kanno, S., Hirano, S., 2001. Oxidative damage to mitochondrial is a preliminary step to caspase-3 activation in fluoride-induced apoptosis in HL60 cells. *Free Rad. Biol. Med.* 31, 367–373.
- Baetu, T.M., Hiscott, J., 2002. On the TRAIL to apoptosis. *Cytokine Growth Factor Rev.* 13, 199–207 (Review).
- Boccellino, M., Pedata, P., Castiglia, L., La Porta, R., Pieri, M., Quagliuolo, L., Acampora, A., Sannolo, N., Miraglia, N., 2010. Doxorubicin can penetrate nitrile gloves and induces apoptosis in keratinocytes cell lines. *Toxicol. Lett.* 197, 61–68.
- Bonfoco, E., Krainc, D., Ankarcona, M., Nicotera, P., Lipton, S.A., 1995. Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with N-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc. Natl. Acad. Sci. U. S. A.* 92, 7162–7166.
- Brenes, O., Muñoz, E., Roldán-Rodríguez, R., Díaz, C., 2010. Cell death induced by *Bothrops asper* snake venom metalloproteinase on endothelial and other cell lines. *Exp. Mol. Pathol.* 88, 424–432.
- Chan, A.B., Allen, C.N., Simmons, N.L., Parsons, M.E., Hirst, B.H., 1989. Resistance to acid of canine kidney (MDCK) and human colonic (T84) and ileo-caecal (HCT-8) adenocarcinoma epithelial cell monolayers in vitro. *Q. J. Exp. Physiol.* 74, 553–556.
- Chen, W.C., Cheng, H.H., Huang, C.J., Chou, C.T., Liu, S.I., Chen, I.S., Hsu, S.S., Chang, H.T., Huang, J.K., Jan, C.R., 2006. Effect of riluzole on Ca²⁺ movement and cytotoxicity in Madin-Darby canine kidney cells. *Hum. Exp. Toxicol.* 25, 461–469.
- Collares-Buzato, C.B., de Paula Le Sueur, L., da Cruz-Hofling, M.A., 2002. Impairment of the cell-to-matrix adhesion and cytotoxicity induced by *Bothrops moojeni* snake venom in cultured renal tubular epithelia. *Toxicol. Appl. Pharmacol.* 181, 124–132.
- Collares-Buzato, C.B., Jepson, M.A., McEwan, G.T., Simmons, N.L., Hirst, B.H., 1994. Junctional uvomorulin/E-cadherin and phosphotyrosine-modified protein content are correlated with paracellular permeability in Madin-Darby canine kidney (MDCK) epithelia. *Histochemistry* 101, 185–194.
- Collares-Buzato, C.B., Jepson, M.A., Simmons, N.L., Hirst, B.H., 1998. Increased tyrosine phosphorylation causes redistribution of adherens junction and tight junction proteins and perturbs paracellular barrier function in MDCK epithelia. *Eur. J. Cell Biol.* 76, 85–92.
- Fan, L.L., Sun, G.P., Wei, W., Wang, Z.G., Ge, L., Fu, W.Z., Wang, H., 2010. Melatonin and doxorubicin synergistically induce cell apoptosis in human hepatoma cell lines. *World J. Gastroenterol.* 16, 1473–1481.
- Ferri, K.F., Kroemer, G., 2001a. Mitochondria—the suicide organelles. *Bioessays* 23, 111–115 (Review).
- Ferri, K.F., Kroemer, G., 2001b. Organelle-specific initiation of cell death pathways. *Nat. Cell Biol.* 3, E255–E263 (Review).
- Gottlieb, E., Vander Heiden, M.G., Thompson, C.B., 2000. Bcl-XL prevents the initial decrease in mitochondrial membrane potential and subsequent reactive oxygen species production during tumor necrosis factor alpha-induced apoptosis. *Mol. Cell Biol.* 20, 5680–5689.
- Jorge, R.J., Monteiro, H.S., Gonçalves-Machado, L., Guarnieri, M.C., Ximenes, R.M., Borges-Nojosa, D.M., Luna, K.P., Zingali, R.B., Corrêa-Netto, C., Gutiérrez, J.M., Sanz, L., Calvete, J.J., Pla, D., 2015. Venomics and antivenomics of *Bothrops erythromelas* from five geographic populations within the Caatinga ecoregion of northeastern Brazil. *J. Proteomics* 114, 93–114.
- Kim, H., Yoon, S.C., Lee, T.Y., Jeong, D., 2009. Discriminative cytotoxicity assessment based on various cellular damages. *Toxicol. Lett.* 184, 13–17.
- Krysko, D.V., Kaczmarek, A., Krysko, O., Heyndrickx, L., Woznicki, J., Bogaert, P., Cauwels, A., Takahashi, N., Magez, S., Bachert, C., Vandennebe, P., 2011. TLR-2 and TLR-9 are sensors of apoptosis in a mouse model of doxorubicin-induced acute inflammation. *Cell Death Differ.* 18, 1316–1325.
- Kusma, J., Chaim, O.M., Wille, A.C., Ferrer, V.P., Sade, Y.B., Donatti, L., Gremski, W., Mangili, O.C., Veiga, S.S., 2008. Nephrotoxicity caused by brown spider venom phospholipase-D (dermonecrotic toxin) depends on catalytic activity. *Biochimie* 90, 1722–1736.
- Landeta, O., Landajuela, A., Gil, D., Taneva, S., DiPrimo, C., Sot, B., Valle, M., Frolov, V.A., Basañez, G., 2011. Reconstitution of proapoptotic BAK function in liposomes reveals a dual role for mitochondrial lipids in the BAK-driven membrane permeabilization process. *J. Biol. Chem.* 286, 8213–8230.
- Marinho, A.D., Morais, I.C., Lima, D.B., Jorge, A.R., Jorge, R.J., Menezes, R.R., Mello, C.P., Pereira, G.J., Silveira, J.A., Toyama, M.H., Orzáez, M., Martins, A.M., Monteiro, H.S., 2015. *Bothropoides pauloensis* venom effects on isolated perfused kidney and cultured renal tubular epithelial cells. *Toxicon* 108, 126–133.
- Martins, M.S., Mendes, M.M., Shimizu, M.H., Melo Rodrigues, V., de Castro, I., Filho, S.R., Malheiros, D.M., Yu, L., Burdman, E.A., 2014. Effects of *Schizolobium parahyba* extract on experimental *Bothrops* venom-induced acute kidney injury. *PLoS One* 9, e86828.
- Martins, A.M.C., Sousa, F.C.M., Barbosa, P.S.F., Toyama, M.H., Toyama, D.O., Aprígio, C.C., Queiroz, M.G.R., Guarnieri, M.C., Havt, A., De Menezes, D.B., Fonteles, M.C., Monteiro, H.S.A., 2005. Action of anti-bothropic factor isolated from *Didelphis marsupialis* on renal effects of *Bothrops erythromelas* venom. *Toxicon* 46, 595–599.
- Martirosyan, A., Clendening, J.W., Goard, C.A., Penn, L.Z., 2010. Lovastatin induces apoptosis of ovarian cancer cells and synergizes with doxorubicin: potential therapeutic relevance. *BMC Cancer* 10, 1–13.
- Masuda, S., Hayashi, H., Araki, S., 1998. Two vascular apoptosis-inducing proteins from snake venom are members of the metalloprotease/disintegrin family. *Eur. J. Biochem.* 253, 36–41.
- Masuda, S., Ohta, T., Kaji, K., Fox, J.W., Hayashi, H., Araki, S., 2000. cDNA cloning and characterization of vascular apoptosis-inducing protein 1. *Biochem. Biophys. Res. Commun.* 278, 197–204.
- Masuda, S., Hayashi, H., Atoda, H., Morita, T., Araki, S., 2001. Purification, cDNA cloning and characterization of the vascular apoptosis-inducing protein, HV1, from *Trimeresurus flavoviridis*. *Eur. J. Biochem.* 268, 3339–3345.
- Mello, C.P., Morais, I.C., Menezes, R.R., Pereira, G.J., Torres, A.F., Lima, D.B., Pereira, T.P., Toyama, M.H., Monteiro, H.S., Smali, S.S., Martins, A.M., 2014. *Bothropoides insularis* venom cytotoxicity in renal tubular epithelia cells. *Toxicon* 88, 107–114.
- Morais, I.C., Torres, A.F., Pereira, G.J., Pereira, T.P., Pessoa Bezerra de Menezes, R.R., Mello, C.P., Coelho Jorge, A.R., Bindá, A.H., Toyama, M.H., Monteiro, H.S., Smali, S.S., Martins, A.M., 2013. *Bothrops leucurus* venom induces nephrotoxicity in the isolated perfused kidney and cultured renal tubular epithelia. *Toxicon* 61, 38–46.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity. *J. Immunol. Methods* 65, 55–63.
- Nascimento, J.M., Franchi Jr., G.C., Nowill, A.E., Collares-Buzato, C.B., Hyslop, S., 2007. Cytoskeletal rearrangement and cell death induced by *Bothrops alternatus* snake venom in cultured Madin-Darby canine kidney cells. *Biochem. Cell Biol.* 85, 591–605.
- Oliveira, F.N., Brito, M.T., Morais, I.C.O., Fook, S.M.L., Albuquerque, H.N., 2010. Accidents caused by *Bothrops* and *Bothropoides* in the State of Paraíba: epidemiological and clinical aspects. *Rev. Soc. Bras. Med. Trop.* 43, 1–6.
- Orrenius, S., McCabe Jr., M.J., Nicotera, P., 1992. Ca²⁺-dependent mechanisms of cytotoxicity and programmed cell death. *Toxicol. Lett.* 64–65, 357–364 (Review).
- Parrish, A.B., Freel, C.D., Kornbluth, S., 2013. Cellular mechanisms controlling caspase activation and function. *Cold Spring Harb. Perspect. Biol.* 5, a008672.
- Peixoto, E.B., Collares-Buzato, C.B., 2005. Protamine-induced epithelial barrier disruption involves rearrangement of cytoskeleton and decreased tight junction-associated protein expression in cultured MDCK strains. *Cell Struct. Funct.* 29, 165–178.
- Queiroz, G.P., Pessoa, L.A., Portaro, F.C.V., Furtado, F.M.D.E., Tambourgi, D.V., 2008. Interspecific variation in venom composition and toxicity of Brazilian snakes from *Bothrops* genus. *Toxicon* 52, 842–851.
- Rocha, I.C.A., 2008. Estudo epidemiológico dos acidentes ofídicos no estado do Ceará no período de 2001 a 2007. Dissertação (Mestrado em Patologia) - Universidade Federal do Ceará, Fortaleza.
- Rodrigues, R.S., Boldrini-França, J., Fonseca, F.P., de la Torre, P., Henrique-Silva, F., Sanz, L., Calvete, J.J., Rodrigues, V.M., 2012. Combined snake venomics and venom gland transcriptomic analysis of *Bothropoides pauloensis*. *J. Proteom.* 75, 2707–2720.
- Sanchez, E.F., Freitas, T.V., Ferreira-Alves, D.L., Velarde, D.T., Diniz, M.R., Cordeiro, M.N., Agostini-Cotta, G., Diniz, C.R., 1992. Biological activities of venoms from South American snakes. *Toxicon* 30, 95–103.
- Schmitz, I., Kirchoff, S., Krammer, P.H., 2000. Regulation of death receptor mediated apoptosis pathways. *Int. J. Biochem. Cell Biol.* 32, 1123–1136 (Review).
- Schwerdt, G., Freudinger, R., Schuster, C., Silbernagl, S., Gekle, M., 2004. Inhibition of mitochondria and extracellular acidification enhance achratoxin A-induced apoptosis in renal collecting duct-derived MDCK-C7 cells. *Cell Physiol. Biochem.* 14, 47–56.
- Sgrignolli, L.R., Mendes, G.E.F., Carlos, C.P., Burdman, E.A., 2011. Acute kidney injury caused by *Bothrops* snake venom. *Nephron Clin. Pract.* 119, 131–137.
- Shamas-Din, A., Kale, J., Leber, B., Andrews, D.W., 2013. Mechanisms of action of Bcl-2 family proteins. *Cold Spring Harb. Perspect. Biol.* 5, a008714.
- Sitprija, V., Sitprija, S., 2012. Renal effects and injury induced by animal toxins. *Toxicon* 5, 943–953.
- Wang, C.J., Zhang, H., Chen, K., Zheng, J.W., Xiao, C.W., Ji, W.W., Yu, Y., Hu, H.Y., Li, Y., Xue, X.B., 2010. Ad.mda-7 (IL-24) selectively induces apoptosis in hepatocellular carcinoma cell lines, suppresses metastasis, and enhances the effect of doxorubicin on xenograft tumors. *Oncol. Res.* 18, 561–574.
- Warrell, D.A., 2004. Snakebites in Central and South America: epidemiology, clinical features and clinical management. In: Campbell, J.A., Lamar, W.W. (Eds.), *Venomous Reptiles of the Western Hemisphere*, vol. 2. Comstock Publishing Associates/Cornell University Press, Ithaca, New York, pp. 709–761.