



Meristiella echinocarpa lectin (MEL): a new member of the OAAH-lectin family

Renata Pinheiro Chaves¹ · Suzete Roberta da Silva² · João Pedro Freire Alves da Silva¹ · Rômulo Farias Carneiro¹ · Bruno Lopes de Sousa³ · Jade Oliveira Abreu⁴ · Fátima Cristiane Teles de Carvalho⁴ · Cintia Renata Costa Rocha⁵ · Wladimir Ronald Lobo Farias¹ · Oscarina Viana de Sousa⁴ · André Luiz Coelho Silva⁶ · Alexandre Holanda Sampaio¹ · Celso Shiniti Nagano¹

Received: 20 February 2018 / Revised and accepted: 4 April 2018 / Published online: 2 May 2018
© Springer Science+Business Media B.V., part of Springer Nature 2018

Abstract

A new lectin from the marine red alga *Meristiella echinocarpa* (MEL) was isolated and biochemically characterized. MEL is a monomeric protein of 28 kDa with specificity for yeast mannan. Hemagglutination activity of MEL was stable between pH 5 and 10, temperatures up to 50 °C, and neither EDTA nor divalent ions affected it. The complete amino acid sequence of MEL was determined through a combination of tandem mass spectrometry and DNA cloning. As a new member of the OAAH-lectin family, the primary structure of MEL consists of 267 amino acid residues distributed in four tandem repeat domains, sharing at least 48% of identity. Theoretical secondary structure of MEL was composed of 3% α -helix, 40% β -sheet, 19% β -turn, and 38% coil. Melting temperatures of the lectin in the absence and presence of mannan were 54 and 61 °C, respectively. Furthermore, MEL was able to recognize and agglutinate pathogenic bacterial strains, such as multidrug-resistant *Salmonella* and *Vibrio alginolyticus*.

Keywords Lectin · Rhodophyta · OAAH-family · Pathogenic bacteria

This work is dedicated in memory of Wladimir Ronald Lobo Farias.

✉ Celso Shiniti Nagano
naganocs@gmail.com

¹ BioMar-Lab, Departamento de Engenharia de Pesca, Universidade Federal do Ceará, Campus do Pici s/n, Av. Mister Hull, bloco 871, Fortaleza, Ceará 60440-970, Brazil

² Campus Monte Alegre, Universidade Federal do Oeste do Pará, Av. Major Francisco Mariano s/n - Bairro Cidade Alta, Monte Alegre, PA 68220-000, Brazil

³ Faculdade de Filosofia Dom Aureliano Matos, Universidade Estadual do Ceará, Av. Dom Aureliano Matos, 2060, Limoeiro do Norte, CE 62930-000, Brazil

⁴ Instituto de Ciências do Mar – Labomar, Universidade Federal do Ceará, Av. da Abolição, 3207, Fortaleza, CE 60165-081, Brazil

⁵ Departamento de Bioquímica, Universidade Federal de Pernambuco, Av. Prof. Moraes Rego, 1235, Cidade Universitária, Recife, PE 50670-901, Brazil

⁶ Departamento de Bioquímica, Universidade Federal do Ceará, Campus do Pici s/n bloco 907, Fortaleza, CE 60440-970, Brazil

Introduction

Lectins are carbohydrate-binding proteins. Although they are found in all organisms, attention has been focused almost exclusively on land plants, whereas other groups, such as marine algae and invertebrates, were often neglected by researchers (Vasta and Ahmed 2008). Interest in algae lectins began 50 years ago. So far, the number of studies with these lectins has grown at a slow pace, mainly owing to the complexity of their isolation caused by the presence of pigments, low concentration of lectins in extracts, creating difficulty in obtaining sufficient material for study, and presence of isoforms in the preparations (Rogers and Hori 1993; Calvete et al. 2000; Nascimento-Neto et al. 2012; Chernikov et al. 2013).

Recently, marine red algae have attracted some attention as potential sources of new lectins with antiviral and anti-HIV activities through binding to viral envelope glycoproteins based on their specificity for high-mannose *N*-glycan (Sato et al. 2011; Hirayama et al. 2016). For instance, griffithsin (GRFT) isolated from *Griffithsia* sp. and KAA isolated from *Kappaphycus alvarezii* showed potent anti-HIV activity (Mori et al. 2005; Hirayama et al. 2016). In addition, other lectins, such as ESA-2 from *Eucheuma serra*, EDA from *Eucheuma denticulatum*, and KSA from *Kappaphycus striatum*, also show affinity for high-mannose *N*-glycans (Hori et al. 2007; Hung et al. 2011, 2015a, b).

With the exception of GFRT, which is a jacalin-like lectin (Mori et al. 2005), most marine red algae lectins with specificity for *N*-glycans have four tandem repeat domains of about 67 amino acids each, showing similarity to a 13-kDa lectin isolated from the cyanobacterium *Oscillatoria agardhii* (OAA). Moreover, these lectins showed identity with lectins isolated from such bacteria as PFL, MBHA, and BOA from *Pseudomonas fluorescens*, *Myxococcus xanthus*, and *Burkholderia oklahomensis*, respectively. OAA was the first lectin of this group to show anti-HIV activity through binding to gp120. Moreover, the tridimensional structure of OAA has been determined (Koharudin et al. 2011); therefore, many authors have termed this group as the OAAH (*O. agardhii* agglutinin homologs) family.

Meristiella echinocarpa (Areschoug) Cheney is a marine red alga found along the northeastern Brazilian coast. Ainouz et al. (1992) showed that *M. echinocarpa* aqueous extract caused high agglutination of rabbit, goat, and chicken erythrocytes. Now, we have isolated and characterized a lectin from *M. echinocarpa* (MEL) with agglutination activity against multidrug-resistant *Salmonella* and *Vibrio alginolyticus*. Furthermore, its primary structure was determined by a combination of MS/MS and DNA cloning.

Material and methods

Collection

Specimens of *Meristiella echinocarpa* were collected in the intertidal zone at Paracuru Beach, Ceará, Brazil. The algae were transported in plastic bags to the lab where they were cleaned from epiphytes, washed with water, and freeze-dried. A small portion of the alga was stored at -80°C for DNA extraction. All collections were authorized through our registration with SISBIO (Sistema de Autorização e Informação em Biodiversidade, ID: 33913-8) and SISGEN (Sistema Nacional de Gestão do Patrimônio Genético e do Conhecimento Tradicional Associado, ID: AC14AF9).

Lectin purification

Freeze-dried algae were triturated until a fine powder was obtained. The algae powder was suspended in twenty volumes of cold ethanol 70% *v/v* and maintained under agitation for 30 min at room temperature. The mixture was then filtered, and solid residues were re-extracted with 70% *v/v* ethanol, as described above, two more times. After that, solid residues recovered by filtration were suspended in ten volumes of 20% *v/v* ethanol and maintained under agitation for 4 h at 4°C . The mixture was filtrated and centrifuged at $8000\times g$ for 20 min at 4°C , and supernatant (crude extract) was stored at -20°C .

Crude extract was dialyzed against sodium phosphate buffer, 20 mM, pH 7 (PB), and loaded onto a DEAE (Diethylaminoethyl)-Sephacel column (1.0×7.0 cm), previously equilibrated with PB. Unbound proteins were washed with equilibrium buffer (fraction D1), and retained proteins were eluted in two steps: PB containing 0.5 M NaCl (fraction D2) and PB containing 1 M NaCl (fraction D3). The chromatography was monitored at 280 nm, 3-mL fractions were collected, and flow rate was maintained at 1.5 mL min^{-1} .

Biochemical properties of MEL

Hemagglutination activity and inhibition by sugars

Hemagglutination activity and inhibition assays were conducted as described by Sampaio et al. (1998), using human (A, B, and O) and rabbit erythrocytes in their native form and treated with the proteases trypsin and papain.

The following sugars and glycoproteins were used in the inhibition assay: D-xylose, D-ribose, L-fucose, L-arabinose, L-rhamnose, D-galactose, D-mannose, D-glucose, D-glucosamine, D-galactosamine, *N*-acetyl-D-glucosamine, *N*-acetyl-D-galactosamine, *N*-acetyl-D-manosamine, D-galacturonic acid, D-fructose, D-sucrose, D-melibiose, α -D-lactose, β -D-lactose, D-lactulose, D-maltose, D-raffinose, mannan from *Saccharomyces cerevisiae* (yeast mannan), methyl- α -D-galactopyranoside, methyl- β -D-galactopyranoside, methyl- β -D-thiogalactose, phenyl- β -D-galactopyranoside, 4-nitrophenyl- α -D-galactopyranoside, 4-nitrophenyl- β -D-galactopyranoside, 2-nitrophenyl- β -D-galactopyranoside, bovine fetuin, and type 2 porcine stomach mucin (PSM).

The effects of pH, temperature, EDTA (ethylenediaminetetraacetic acid) and divalent cations on hemagglutination activity were evaluated, as described in pre-established methods (Sampaio et al. 1998).

Carbohydrate content

Neutral carbohydrate content in MEL was estimated as described by Dubois et al. (1956), using lactose as standard.

Molecular mass determination

Homogeneity and molecular mass of MEL were estimated by SDS-PAGE (Laemmli 1970) in the presence and absence of 2-mercaptoethanol (2-ME). LMW-SDS Marker kit (GE Healthcare, UK) was used as molecular weight marker.

Native molecular mass of MEL was estimated by gel filtration chromatography in BioSuite 250, using a $5\text{-}\mu\text{m}$ HR column coupled to an Acquity UPLC system (Waters Corp, USA). Chromatography was performed in sodium phosphate buffer 20 mM, pH 7, containing 0.15 M NaCl (PBS) at a flow rate of 0.4 mL min^{-1} . The column was previously calibrated

with a gel filtration marker kit for protein weights between 29 and 700 kDa (Sigma-Aldrich, USA).

Average molecular mass of MEL was determined by MALDI-ToF on an Autoflex III mass spectrometer (Bruker Daltonics, Germany), using matrix solution (10 mg mL⁻¹ of CHCA (α -cyano-4-hydroxycinnamic acid) acetonitrile, water, and TFA (trifluoroacetic acid), 50:47:3% v/v). The spectra were acquired in linear positive mode and processed with Flex Analysis 3.4 software (Bruker Daltonics, Germany).

Primary structure determination

Tandem mass spectrometry (MS/MS)

First, MEL was subjected to 1D-SDS PAGE as described above. Protein spots were digested with trypsin and chymotrypsin, and peptides were extracted from gel according to Shevchenko et al. (2006). Peptides were separated on a reverse phase C-18 nanocolumn (0.075 × 100 mm) coupled to a nanoAcquity system. The eluates were directly infused in a hybrid mass spectrometer (ESI-Q-ToF) (Synapt HDMS, Waters Corp, USA). The instruments' parameters were adjusted as described by Carneiro et al. (2013). MS/MS spectra were manually interpreted, and sequenced peptides were searched online against NCBI and UniProt databanks.

DNA extraction and purification

Small pieces of algae were ground in dry ice until a fine powder was obtained. Powdered alga was incubated with CTAB extraction buffer (2% cetyl trimethylammonium bromide, 1.4 M NaCl, 0.1 M Tris-HCl, 20 mM EDTA, and 2% 2-ME; Sigma-Aldrich) containing 1% PVPP (polyvinylpyrrolidone) in water bath at 60 °C for 1 h. The extract was centrifuged at 9000×g for 20 min, and the supernatant was transferred to a clean tube. One volume of chloroform/isoamyl alcohol solution

(24:1) was added, and the mixture was centrifuged at 9000×g for 20 min. The upper phase was collected, washed with chloroform/isoamyl alcohol solution, centrifuged, and transferred to a clean tube. A 0.3 volume of isopropyl alcohol was added; then, the mixture was incubated for 20 min at -20 °C and centrifuged at 9000×g for 10 min. The supernatant was removed, and 1 mL of 70% cold ethanol was added to the precipitate. Again, the mixture was centrifuged at 9000×g for 10 min, and the supernatant was removed. The precipitate was dried at room temperature and then rehydrated with a small volume of nuclease-free water.

Primers, PCR, and product sequencing

Degenerate primers were designed based on the amino acid sequences obtained by MS/MS: VQNQWGG (primer-Upst-VQ-FW: 5'- GTI CAG AAT CAR TGG GGI GG -3') and EGPIGF (primer-Down-Rv: 5'- RAA GCC GAT BGG WCC TTC -3'). PCR was performed with a 25- μ L reaction mixture containing 100 ng of the genomic DNA, 0.4 μ M of each primer, 0.4 mM of dNTP Mix (Promega, USA), and 1 U of Platinum *Taq* DNA polymerase (Invitrogen, USA) in 1X PCR buffer with 3 mM of MgCl₂. The amplification protocol included an initial denaturation step at 94 °C for 5 min, followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 50 °C for 60 s, extension at 72 °C for 60 s, and the final extension step at 72 °C for 5 min.

The PCR product was purified by the PureLink Quick Gel Extraction Kit (Invitrogen). Then, the PCR product was cloned into pGEM-T Easy Vector, transformed into *Escherichia coli* strain DH5 α (Novagen, Brasil), and screened with blue-white selection in LB agar containing 100 μ g mL⁻¹ ampicillin (Thermo Scientific, USA), 0.5 mM IPTG (isopropyl β -D-1-thiogalactopyranoside; Thermo Scientific), and 80 μ g mL⁻¹ of X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside; Thermo Scientific). The cloning was

Fig. 1 Ion exchange chromatography. Approximately 40 mL of dialyzed extract were applied onto a DEAE-Sephacel column equilibrated with phosphate buffer, pH 7. The column was washed with the equilibrium buffer, and two adsorbed fractions (D2 and D3) were eluted with 0.5 and 1 M of NaCl in the phosphate buffer




Table 1 Summary of purification of MEL

Fraction	Protein concentration (mg mL ⁻¹)	Total activity (HU mL ⁻¹)	Specific activity (HU mg ⁻¹)	Purification (fold)
Crude extract	2.080	5120	2.462	1.0
D2	0.244	3840	15.754	6.4

performed in a biosafety laboratory certified in accordance with governmental requirements (CQB: R007-2016).

Recombinant plasmids were extracted by the illustra plasmid Prep Mini Spin Kit (GE Healthcare, UK) and confirmed by restriction digestion with EcoRI (Promega). Finally, constructions were sequenced in the MegaBACE automatic sequencer (GE Healthcare), using primers containing the T7 (5'-TAA TAC GAC TCA CTA TAG GG-3') and SP6 promoter (5'-ATT TAG GTG ACA CTA TAG-3') sequences.

Sequencing was performed with at least six clones from PCR cloning amplification to avoid any PCR errors. The reads were analyzed by the Phred-Phrap-Consed program. The contigs formed were translated into amino acids, using the ExPASy translation tool (<https://web.expasy.org/translate/>).

Circular dichroism

MEL (0.2 mg mL⁻¹ in 20 mM phosphate buffer, pH 7.0, containing 100 mM NaCl) was placed in a rectangular quartz cuvette with 0.5 mm path length. Spectra were acquired at a scan speed of 50 nm min⁻¹ with a bandwidth of 1 nm in a Jasco J-815 spectropolarimeter (Jasco International Co., Japan) connected to a peltier module with controlled temperature. The acquisitions were performed at 190–250 nm (far UV). The DICHROWEB web server (Whitmore and Wallace 2008) was used to perform analyses of secondary structure prediction.

The thermodynamic parameters of lectin folding and unfolding in the presence and absence of ligands were calculated by monitoring the changes in ellipticity at 214 nm as a function of temperature (Greenfield 2007). The lectin was evaluated in the presence and absence of yeast mannan (0.1 mg mL⁻¹).

Antibacterial activity and agglutination assays

The following standard strains were used in tests: *Escherichia coli* ATCC 25922 and *Staphylococcus aureus* ATCC 25923, multidrug-resistant *Salmonella* ser. Brandeirup Lamap 18 (Lamap collection), *Vibrio cholerae* IOC 19582, *V. parahaemolyticus* IOC 18950, *V. harveyi* ATCC 14126, and *V. alginolyticus* ATCC 17749.

Antibacterial activity of MEL was performed in microtiter plates according to the broth dilution method as described by Balouiri et al. (2016). Bacterial inocula were prepared in tubes containing 0.85 and 1% NaCl solutions. Turbidity was adjusted to 0.5 according to the McFarland scale. Each well containing Mueller-Hinton broth received 50 µL of bacterial inoculum and

50 µL of MEL at 500, 250, 100, and 50 µg mL⁻¹. Kanamycin A (Affymetrix USB products, USA) at 30 µg mL⁻¹ was used as positive control, and bacterial inoculum without lectin was used as negative control. The procedure was performed in triplicate. The plates were incubated at 28 °C for 48 h, and then triphenyl tetrazolium chloride (TTC) (Sigma-Aldrich) was added at the

Table 2 Inhibitory effects of sugars and glycoproteins in hemagglutinating activity of MEL. The initial concentrations of the inhibitors were 100 mM for sugars and 1 mg mL⁻¹ for glycoproteins

Sugar	MIC
D-galactose	NI
D-glucose	NI
D-mannose	NI
D-xylose	NI
D-ribose	NI
L-fucose	NI
L-rhamnose	NI
L-arabinose	NI
α-Methyl-D-galactopyranoside	NI
β-Methyl-D-galactopyranoside	NI
Methyl-β-D-thiogalactose	NI
D-galactosamine	NI
D-glucosamine	NI
D-N-acetyl-galactosamine	NI
D-N-acetyl-glucosamine	NI
D-N-acetyl-manosamine	NI
D-galacturonic acid	NI
Phenyl-β-D-galactopyranoside	NI
2-nitro-phenyl-β-D-galactoside	NI
4-nitro-phenyl-α-D-galactoside	NI
4-nitro-phenyl-β-D-galactoside	NI
D-fructose	NI
D-saccharose	NI
α-Lactose	NI
β-Lactose	NI
D-maltose	NI
Lactulose	NI
Melibiose	NI
Raffinose	NI
Yeast mannan	62.5 µg mL ⁻¹
Glycoprotein	
PSM type II	NI
Bovine fetuin	NI

MIC minimal inhibitory concentration, NI inhibition not observed

final concentration of 0.5%. The reaction was incubated at 28 °C for 3 h. The change of color to red was considered negative activity.

The agglutination assay was conducted according to Melo et al. (2014). The lectin ($100 \mu\text{g mL}^{-1}$) was incubated with bacteria in the presence and absence of yeast mannan at $100 \mu\text{g mL}^{-1}$. Results were observed after incubation for 1 h under a light microscope.

Results

Lectin purification

The crude extract of *M. echinocarpa* showed hemagglutination activity against trypsin-treated rabbit erythrocytes. After ion exchange chromatography, hemagglutination activity was concentrated in fraction D2. Fractions D1 and D3 showed residual activity (Fig. 1). The purification procedure increased by 6.4-fold the lectin activity compared to the crude extract, and MEL represented 75% of the total hemagglutination activity present in the crude extract (Table 1).

Biochemical properties of MEL

MEL could agglutinate only rabbit erythrocytes treated with papain (data not shown).

Simple sugars could not inhibit hemagglutination activity caused by MEL. However, the lectin was inhibited by yeast

mannan with minimal inhibitory concentration (MIC) of $62.5 \mu\text{g mL}^{-1}$ (Table 2). Hemagglutination activity of MEL was stable between pH 5 and 10. Lectin activity was unaltered up to 50 °C, but after that, activity was gradually lost and then completely abolished at 70 °C (data not shown). The presence of EDTA, CaCl_2 , MnCl_2 , and MgCl_2 did not affect hemagglutination activity caused by MEL.

The phenol-sulfuric acid method suggested that MEL is a glycoprotein with less than 2% of neutral sugar in its composition. In SDS-PAGE, MEL showed a broad band of 30 kDa in the presence and absence of 2-ME (Fig. 2). Under native conditions (gel filtration chromatography), MEL was determined to be a monomeric protein with relative molecular mass (Mr) of 21 kDa (data not shown).

MALDI-ToF analysis revealed a broad signal between 28,000 and 30,000 Da with maximal intensity at 28,907 Da (Fig. 2).

Amino acid sequence determination

The primary structure of MEL was determined by a combination of mass spectrometry and DNA cloning. The amplification product obtained by DNA cloning was approximately 400 bp. This DNA sequence encodes a polypeptide chain of 124 amino acids, which corresponds to an amino acid sequence between 4 and 127 residues. The amino acid sequences of peptides obtained by digestion with trypsin and chymotrypsin were sufficient to cover of the primary structure of MEL, in addition to confirming the deduced sequence obtained by DNA (Fig. 3).




Fig. 2 Molecular mass determination. MALDI-ToF spectra of MEL. Insert. Electrophoresis profile of purified MEL. Purified MEL in the absence and presence of 2-mercaptoethanol (lanes 1 and 2, respectively). M, molecular marker

Fig. 3 Primary structure of MEL. Amino acid sequence of MEL assembled from sequences of overlapping peptides generated by cleavage with trypsin (-T) and chymotrypsin (-Q) and deduced DNA sequences (underlined)

GRYTVQNWQG GSSAPWNDAG LYVLGGRANQ NVVAIEVSSN DGGADLSGTM TYSGEGPIGF KGTRRGESNV
 |-----Q1-----| |-----T2-----|
 |-----T1-----| |-----Q2--|
 HEVKNQWGGG SAPWHDGGDF VIGSRSGQGV VGLDIKSSDN GKTLTGTMTY EREGPIGFKG TQSGGDTYNV
 |-----T3-----| |-----T4-----| |-----T5-----| |-----T6-----| |-----T7-----|
 |-----Q3-----| |-----Q4-----| |-----Q5-----|
 ENQWGGSSAP WNKAGVWALG DRSSSEAMNKL EVSSSDGGST LEGTMQYKGE GPIGFRAKLA SANNYSVENQ
 -----T7-----| |-----T8-----| |-----T9-----| |-----T10-----|
 ---Q5-----| |-----Q7-----| |-----Q8-----| |-----Q9-----| |-----Q10-----|
 |-----Q11-----| |-----T11-----| |-----T12-----| |-----T13-----| |-----T14-----|
 |-----Q12-----| |-----Q13-----|

	1	10	20	30	40	50	60
MEL	.GRYTVQNWQGGSSAPWNDAGLYVVLGGRANQNVVAIEVSSNDGGADLSGTMTYSGEGPIGFKGTRRGESN						
MPA-2	MGRYTVQNWQGGSSAPWNDAGLYVVLGSRANQNVVAIEVSSNDGGANLGTMTYSGEGPIGFKGARRGESN						
MPA-1	MGRYTVQNWQGGSSAPWNDAGLYVVLGSRANQNVVAIEVSSNDGGANLGTMTYSGEGPIGFKGARRGQSN						
KAA-1	MGRYTVQNWQGGSSAPWNDAGLWILGSRGNQNVMAIDVNSSDGGANLNGTMTYSGEGPIGFKGARRGESN						
KSA-2	MGRYTVQNWQGGSSAPWNDAGLWILGSRGNQNVMAIDVNSSDGGANLNGTMTYSGEGPIGFKGARRGESN						
KAA-2	MGRYTVQNWQGGSSAPWNDAGLWILGSRGNQNVMAIDVNSSDGGANLNGTMTYSGEGPIGFKGARRGDSN						
ESA-2	.GRYTVQNWQGGSSAPWNDAGLWILGSRGNQNVMAIDVNSSDGGANLNGTMTYSGEGPIGFKGARRGESN						
EDA-2	MGRYTVQNWQGGSSAPWNDAGLCILGSRGNQNVIAVDVTSDDGGANLGTMTYSGEGPIGFKGARRGESN						
ASL-1	MGRYTVQNWQGGSSAPWNEAGLWVLGGRANQNVMAIDVSSDGGANLGTMTYSGEGPIGFKGTRRGNSN						
ASL-2	MARYTVQNWQGGSSAPWNDAGLYILGNRANQNVMAIDVTSDDGGAVLGTMTYSGEGPIGFKGTRRGDST						
consensus>50	mgRYTVQNWQGGSSAPWN#AGLw!LGsRaNQNvMa!#V.SsDGGANL.GTMTYSGEGPIGFKGaRRG#SN						

	70	80	90	100	110	120	130
MEL	VHEVKNQWGGSSAPWHDGGDFVIGSRSGQGVVGLDIKSSDNKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
MPA-2	VYEVKNQWGGSSAPWHDGGDFVIGSRSGQGVVGLDIKSSDNKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
MPA-1	VYEVKNQWGGSSAPWHDGGDFVIGSRSGQGVVGLDIKSSDNKTKLTGTMTYERKGPFGFKGTQSGGDSYN						
KAA-1	VYDVENQWGGSSAPWHAAGGQFVIGSRSGQGVLAVNITSSDGKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
KSA-2	VYDVENQWGGSSAPWHAAGGQFVIGSRSGQGVLAVNITSSDGKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
KAA-2	VYDVENQWGGSSAPWHAAGGQFVIGSRSGQGVLAVNITSSDGKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
ESA-2	VYDVENQWGGSSAPWHAAGGQFVIGSRSGQGVLAVNITSSDGKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
EDA-2	VYDVENQWGGSSAPWHAAGGQFVIGSRSGQGVLAVNITSSDGKTKLTGTMTYEREGPIGFKGTQSGGDTYN						
ASL-1	VYEVENQWGGSSAPWHDGGDFVIGSRSGQGVVGLNVSDDNKGTKLTGTMTYEREGPIGFKGTQSGGDSYN						
ASL-2	VYEVENQWGGSSAPWRPAGGDFVIGSRAAGQGVVGLDVSDDNKGTKLTGTMTYEREGPIGFKGTQSGGDSYN						
consensus>50	Vy#VeNQWGGSSAPWHaGGqFVIGSRsGQGVvaln!tSSDNKTKLTGTMTYEReGPiGFKGTQSGGDTYN						

	140	150	160	170	180	190	200
MEL	VENQWGGSSAPWNKAGVWALGDRSSSEAMNKLKLEVSSSDGGSTLEGTMOYKGEPIGFRAKLASANNYSVEN						
MPA-2	VENQWGGSSAPWNKAGVWALGDRSSQAMIKLEVSSSDGGSTLEGTMOYKGEPIGFRAKLASANNYSVEN						
MPA-1	VENQWGGSSAPWNKAGI WALGDRAGQAMIKLEVTSDDGGSNLEGTMOYKGEPIGFRAKLSGANNYSVEN						
KAA-1	VENQWGGSSAPWNKAGI WALGDRSGQAMIAMDVSSSDGGKTLLEGTMOYKGEPIGFRGKLSGANNYSVEN						
KSA-2	VENQWGGSSAPWNKAGI WALGDRSGQAMIAMDVSSSDGGKTLLEGTMOYKGEPIGFRGKLSGANNYSVEN						
KAA-2	VENQWGGSSAPWNKAGI WALGDRSGQAMIAMDVSSSDGGKTLLEGTMOYKGEPIGFRGKLSGANNYSVEN						
ESA-2	VENQWGGSSAPWNKAGI WALGDRSGQAMIAMDVSSSDGGKTLLEGTMOYKGEPIGFRGKLSGANNYSVEN						
EDA-2	VENQWGGSSAPWNKAGI WALGDRNGQAMIAMDVSSSDGGKTLLEGTMOYKGEPIGFRGKLSGANNYSVEN						
ASL-1	VENQWGGSSAAWNKAGVWALGDRNGQGVIGVDVTSDDGGKTLLEGTMOYKGEPIGFRGKLSANNYSVEN						
ASL-2	VENQWGGSSAPWNKAGI WALGDRNGQGMIGLDVTSDDGGKTLLEGTMOYKNEGPIGFRGKLSGANNYSVEN						
consensus>50	VENQWGGSSApWNKAG!WALGDRsg#amiam#VsSsDGGktLEGTmOYKgeGPIGFRgKLSgANNYSVEN						

	210	220	230	240	250	260
MEL	QWGGSSAPWDKAGDWLIGDRYNQNI TAVNVSSSDDGKNGLEGTCTYAREGPIGFKGSAV.					
MPA-2	QWGGSSAPWNKAGDWLIGDRYNQNI TAVNVSSSDDGKNGLEGTCTYAREGPIGFKGSAV.					
MPA-1	QWGGSSAPWNKAGDWLIGDRYNQNI TAVNVASDDGKNGLEGTCTYAREGPIGFKGAAV.					
KAA-1	QWGGSSAPWNKAGDWLIGDRHNQNI TAVKVSDDNDGKNLDGTCTYEREGPIGFKGVAS.					
KSA-2	QWGGSSAPWNKAGDWLIGDRHNQNI TAVKVSDDNDGKNLDGTCTYEREGPIGFKGVATS					
KAA-2	QWGGSSAPWNKAGDWLIGDRHNQNI TAVKVSDDNDGKNLDGTCTYEREGPIGFKGVATS					
ESA-2	QWGGSSAPWNAAGDWLIGDRHNQNI TAVKVSDDNDGKNLDGTCTYEREGPIGFKGVATS					
EDA-2	QWGGSSAPWNKAGDWLIGDRYNQNI TAVKVSDDNDGKNLDGTCTYEREGPIGFKGVATS					
ASL-1	QWGGSSAPWNEAGNWLIGDRHNQNI VALKVTSDDDGKNGLEGTCTYAREGPIGFKGVNS					
ASL-2	QWGGSSAPWNKAGNWLIGDRHNQNI VAVKVSSTDNKGKNGLEGTCTYAREGPIGFKGVAN.					
consensus>50	QWGGSSAPW#kAG#WLIGDRhNQNI tAvkVsSd##GKNL#GTCTYerEGP!GFKGva.s					

Fig. 4 Alignment of MEL and OAAH-lectin family. Alignment obtained by comparison of MEL, *Meristotheca papulosa* agglutinins (MPA-1 and -2), *Agardhiella subulata* lectins (ASL-1 and -2), *Kappaphycus alvarezii*

agglutinins (KAA-1 and -2), *K. striatum* agglutinins (KSA-2), *Euclumea serra* agglutinin (ESA), *E. denticulatum* agglutinin (EDA), and *Solieria filiformis* lectins (SfL-1 and -2). ESPrpt 3.0 was used as alignment tool

The primary structure of MEL consisted of 267 residues, including one cysteine, which did not appear to be involved in disulfide bond formation. Three *N*-glycosylation sequences were found along the sequence, but no carbohydrates were found attached to the polypeptide backbone during MS or MS/MS analysis. Theoretical pI and Mr of MEL were 4.9 and 27,942 Da, respectively. The amino acid sequence of MEL was similar to several members of the OAAH-family (Fig. 4), including MPAs from *Meristotheca papulosa* [BAX08602.1; BAX08602.1], ASLs from *Agardhiella subulata* [BAX08599.1; BAX08598.1], KAA [BAU19431.1; BAU19430.1], KSA-2 [BAR91206.1], ESA-2 [P84331.1], EDA-2 [BAR91516.1], and SfL-1 and -2 from *Solieria filiformis* (Chaves et al. 2018).

Circular dichroism

Native MEL exhibited minimum absorption at 207 and 217 nm in CD measurements (Fig. 5). As determined by the CONTIN prediction method, a general-purpose constrained regularization method for continuous distributions (Van Stokkum et al. 1990), the theoretical secondary structure of MEL was composed of 3% α -helix, 40% β -sheet, 19% β -turn, and 38% coil.

The protein-ligand complex (MEL-mannan) showed greater resistance to thermal denaturation than the lectin in the absence of ligand (Fig. 6). Melting temperatures (T_m) of the lectin in the absence and presence of mannan were 54 and 61 °C, respectively.

Antibacterial and agglutination assays

MEL showed no antibacterial activity against strains tested. However, MEL was able to agglutinate cells of *V. alginolyticus* and multidrug-resistant *Salmonella*. Agglutinations were not observed when the lectin was previously incubated with yeast




Fig. 5 CD spectra of MEL. Far-UV CD spectra (200–240 nm) of MEL. The cuvette path length was 0.05 cm; protein concentration was 0.2 mg mL⁻¹ in PBS, pH 7

mannan (Fig. 7). MEL showed no agglutination activity against *E. coli*, *S. aureus*, *V. cholerae*, *V. parahaemolyticus*, or *V. harveyi*.

Discussion


A new lectin isolated from the marine red alga *M. echinocarpa* (MEL) showed several characteristics common to other marine red algae lectins, such as Ca²⁺-independent activity, no inhibition for simple sugars, and relative stability to pH and temperature variation (Hori et al. 1990; Rogers and Hori 1993).

In marine red algae, at least three families of lectins can be considered. The first family is composed of lectins isolated from *Hypnea japonica* (HJA) and lectins from the genus *Bryothamnion*, BTL and BSL from *B. triquetrum* and *B. seaforthii*, respectively. These lectins show low molecular weight (~9 kDa), four conserved cysteines, specificity for complex carbohydrates, and similarity of sequence (Ainouz et al. 1995; Calvete et al. 2000; Hori et al. 2000; Nascimento-Neto et al. 2012). The second family includes lectins isolated from *Hypnea musciformis* (HML) and *H. cervicornis* (HCA), which, together, present 14 conserved cysteines, a complex post-translational processing with proteolytic cleavage and rebinding of chains by disulfide bond, and specificity for fucose cores (Nagano et al. 2002, 2005). Finally, the third family, OAAH, is represented by lectins with primary structure similar to OAA. Our finding strongly indicated that MEL could be grouped into this family. MEL showed identity with MPA-2 (97%), MPA-1 (93%), ASL-1 (84%), ASL-2 (83%), KAA-2 (84%), KSA-2 (84%), EDA-2 (84%), SfL-1 (78%), and SfL-2 (78%). Moreover, like other OAAH-family members, MEL has four tandem repeat domains, which shared at least 48% of identity. Interestingly, two short sequences are present in all domains: “NQWGGSSAPW” and “EGPIGF.” These sequences are also present in BOA and correspond to carbohydrate-binding regions (Whitley et al. 2013).

Sequence “EGPIGF” was chosen to design primers in the cloning step. However, owing to the repetition of this sequence in the four domains, several amplification products of varied lengths were obtained after PCR. Curiously, a fragment containing four domains was not observed among the amplification products. Therefore, MS/MS was applied to complete the primary structure of MEL, and results revealed the remainder of the sequence and confirmed the deduced DNA sequence.

MEL showed specificity similar to other OAAH-family members. Like MEL, *Eucheuma* and *Kappaphycus* lectins were inhibited by small amounts of yeast mannan (Kawakubo et al. 1999; Hung et al. 2011; Hirayama et al. 2016). Lectins from the OAAH-family exclusively recognize high-mannose *N*-glycans. Yeast mannan is a highly branched oligomannoside with $\alpha(1 \rightarrow 2)$ - and $\alpha(1 \rightarrow 3)$ -linked side

Fig. 6 Determination of MEL T_m from changes in CD as a function of temperature. MEL solubilized in PBS, pH 7, in the absence (a) and presence (b) of yeast mannan



chains attached to an $\alpha(1 \rightarrow 6)$ -linked backbone (Jones and Ballou 1969). Lectins from the OAAH-family have demonstrated interesting biological properties. For instance, ESA induced cell death against several cancer cell lines, such as colon cancer Colo201 cells and cervix cancer HeLa cells (Sugahara et al. 2001). Sfls showed in vitro effect against human breast cancer (Chaves et al. 2018), and KAA-2 inhibited infection of various influenza strains with EC₅₀ of nanomolar levels (Sato et al. 2011).


ESA-2 and EDA-2 were able to inhibit growth of *Vibrio vulnificus* and *V. alginolyticus*, respectively (Liao et al. 2003; Hung et al. 2015a). EDA activity against *V. alginolyticus* was inhibited by yeast mannan. Similarly, MEL was able to recognize *V. alginolyticus*, but no agglutination effect was observed in the presence of yeast mannan, suggesting that mannoside, or mannoside-like structure(s), must be found on the bacterial surface, acting as receptors for lectins (Hung et al. 2015a). Additionally, the inhibition of MEL by mannan in the bacterial agglutination of multidrug-resistant *Salmonella* also

suggested the presence of mannoside, or mannoside-like, structure(s) on its cell surface.

The bacterial agglutination displayed by some lectins resides in the recognition and interaction with polysaccharide or lipopolysaccharide on the bacterial cell surface, and this ability can be used to identify and distinguish microorganisms according to the different compounds exposed on their cell surfaces. Thus, lectins can be an alternative method for typing microorganisms with minimum specialized facilities. In particular, the agglutination assay is a rapid, inexpensive, reproducible, and simple assay to perform and a useful method for epidemiological studies and detection of bacteria (Ottenssooser et al. 1974; Schaefer et al. 1979; Davidson et al. 1982; Khin et al. 2000; Templier et al. 2016).

In conclusion, we have isolated a new lectin from the OAA-family 25 years after the initial report of Ainouz et al. (1992), who demonstrated hemagglutination activity in aqueous extracts of the *M. echinocarpa*. MEL showed the ability to select and type pathogenic microorganisms.

Fig. 7 Agglutination of bacteria by MEL. *V. alginolyticus* incubated with NaCl 1% (a), MEL (b), and MEL complexed with yeast mannan (c); *Salmonella* incubated with NaCl 0.85% (d), MEL (e), and MEL complexed with yeast mannan (f)



Acknowledgments This work was supported by the Brazilian agencies CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), FUNCAP (Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico), and FINEP (Financiadora de Estudos e Projetos). The authors thank CETENE by the availability of the mass spectrometer, and, especially, the authors are grateful to Dr. Julia Campos for MALDI-ToF experiments. AHS, CSN, and OVS are senior investigators of CNPq.

References

- Ainouz IL, Sampaio AH, Benevides NMB, Feritas ALP, Costa FHF, Carvalho MR, Pinheiro-Joventino F (1992) Agglutination of enzyme treated erythrocytes by Brazilian marine algal extract. *Bot Mar* 35:475–479
- Ainouz IL, Sampaio AH, Freitas ALP, Benevides NMB, Mapurunga S (1995) Comparative study on hemagglutinins from the red algae *Bryothamnion seaforthii* and *Bryothamnion triquetrum*. *R Bras Fisiol Veg* 7:15–19
- Balouiri MM, Sadiki M, Ibsouda SK (2016) Methods for in vitro evaluating antimicrobial activity: a review. *J Pharm Anal* 6:71–79
- Calvete JJ, Costa FHF, Saker-Sampaio S, Murciano MPM, Nagano CS, Cavada BS, Grangeiro TB, Ramos MV, Jr CB, Silveira SB, Freitas BT, Sampaio AH (2000) The amino acid sequence of the agglutinin isolated from the red marine algae *Bryothamnion triquetrum* defines a novel lectin structure. *Cell Mol Life Sci* 57:343–350
- Carneiro RF, Melo AA, Almeida AS, Moura RM, Chaves RP, Sousa BL, Nascimento KS, Sampaio SS, Lima JP, Cavada BS, Nagano CS, Sampaio AH (2013) H-3, a new lectin from the marine sponge *Haliclona caerulea*: purification and mass spectrometric characterization. *Int J Biochem Cell Biol* 45:2864–2873
- Chaves RP, da Silva SR, Nascimento Neto LG, Carneiro RF, Coelho da Silva AL, Sampaio AH, Lopes de Sousa B, Cabral MG, Videira PA, Teixeira EH, Nagano CS (2018) Structural characterization of two isolectins from the marine red alga *Solieria filiformis* (Kützinger) P.W. Gabrielson and their anticancer effect on MCF-7 breast cancer cells. *Int J Biol Macromol* 107(Pt A):1320–1329
- Chernikov OV, Molchanova VI, Chikalovets IV, Kondrashina AS, Li W, Lukyanov PA (2013) Lectins of marine hydrobionts. *Biochem Mosc* 78:760–770
- Davidson SK, Keller KF, Doyle RJ (1982) Differentiation of coagulase-positive and coagulase-negative staphylococci by lectins and plant agglutinins. *J Clin Microbiol* 15:547–553
- Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F (1956) Colorimetric method for determination of sugars and related substances. *Anal Chem* 28:350–356
- Greenfield NJ (2007) Determination of the folding of proteins as a function of denaturants, osmolytes or ligands using circular dichroism. *Nat Protoc* 1:2733–2741
- Hirayama M, Shibata H, Imamura K, Takemasa Sakaguchi T, Hori K (2016) High-mannose specific lectin and its recombinants from a carrageenophyta *Kappaphycus alvarezii* represent a potent anti-HIV activity through high-affinity binding to the viral envelope glycoprotein gp120. *Mar Biotechnol* 18:144–160
- Hori K, Miyazawa K, Ito K (1990) Some common properties of lectins from marine algae. *Hydrobiologia* 204/205:561–566
- Hori K, Matsubara K, Miyazawa K (2000) Primary structures of two hemagglutinins from the marine red alga, *Hypnea japonica*. *Biochim Biophys Acta* 1474:226–236
- Hori K, Sato Y, Ito K, Fujiwara Y, Iwamoto Y, Makino H, Kawakubo A (2007) Strict specificity for high-mannose type N-glycans and primary structure of a red alga *Eucheuma serra* lectin. *Glycobiology* 17:479–491
- Hung LD, Sato Y, Hori K (2011) High-mannose N-glycan-specific lectin from the red alga *Kappaphycus striatum* (carrageenophyte). *Phytochemistry* 72:855–861
- Hung LD, Hirayama M, Ly BM, Hori K (2015a) Purification, primary structure, and biological activity of the high-mannose N-glycanspecific lectin from cultivated *Eucheuma denticulatum*. *J Appl Phycol* 27:1657–1669
- Hung LD, Hirayama M, Ly BM, Hori K (2015b) Biological activity, cDNA cloning and primary structure of lectin KSA-2 from the cultivated red alga *Kappaphycus striatum* (Schmitz) Doty ex Silva. *Phytochem Lett* 14:99–105
- Jones GH, Ballou CE (1969) Studies on the structure of yeast mannan. *J Biol Chem* 244:1043–1051
- Kawakubo A, Makino H, Ohnishi J, Hirohara H, Kanji H (1999) Occurrence of highly yielded lectins homologous within genus *Eucheuma*. *J Appl Phycol* 11:149–156
- Khin MM, Hua JS, Ng HC, Wadström T, Ho B (2000) Agglutination of *Helicobacter pylori* coccoids by lectins. *World J Gastroenterol* 6:202–209
- Koharudin LMI, Furey W, Gronenborn AM (2011) Novel fold and carbohydrate specificity of the potent anti-HIV cyanobacterial lectin. *J Biol Chem* 286:1588–1597
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227:680–683
- Liao WR, Lin JY, Shieh WY, Jeng WL, Huang R (2003) Antibiotic activity of lectins from marine algae against marine vibrios. *J Ind Microbiol Biotechnol* 30:433–439
- Melo AA, Carneiro RF, Silva WM, Moura RM, Silva GC, Sousa OV, Saboya JS, Nascimento KS, Saker-Sampaio S, Nagano CS, Cavada BS, Sampaio AH (2014) HGA-2, a novel galactoside-binding lectin from the sea cucumber *Holothuria grisea* binds to bacterial cells. *Int J Biol Macromol* 64:435–442
- Mori T, O’Keefe BR, Sowder RC, Bringans S, Gardella R, Berg S, Cochran P, Turpin JA, Buckheit RW, McMahon JB, Boyd MR (2005) Isolation and characterization of griffithsin, a novel HIV inactivating protein, from the red alga *Griffithsia* sp. *J Biol Chem* 280:9345–9353
- Nagano CS, Moreno FBMB, Bloch C, Prates MV, Calvete JJ, Sampaio SS, Farias WRL, Tavares TD, Nascimento KS, Grangeiro TB, Cavada BS, Sampaio AH (2002) Purification and characterization of a new lectin from red marine alga *Hypnea musciformis*. *Protein Pept Lett* 9:159–165
- Nagano CS, del Sol FG, Cavada BS, Nascimento KSD, Nunes EV, Sampaio AH, Calvete JJ (2005) Crystallization and preliminary X-ray diffraction analysis of HML, a lectin from the red marine alga *Hypnea musciformis*. *Acta Cryst F* 61:997–999
- Nascimento-Neto LG, Carneiro RF, da Silva SR, da Silva BR, Arruda FVS, Carneiro VA, do Nascimento KS, Saker-Sampaio S, da Silva VA Jr, Porto ALF, Cavada BS, Sampaio AH, Teixeira EH, Nagano CS (2012) Characterization of isoforms of the lectin isolated from the red algae *Bryothamnion seaforthii* and its pro-healing effect. *Mar Drugs* 10:1936–1954
- Ottensoufer F, Nakamizo Y, Sato M, Miyamoto Y, Takizawa K (1974) Lectins detecting group C streptococci. *Infect Immun* 9:971–973
- Rogers DJ, Hori K (1993) Marine algal lectins: new developments. *Hydrobiologia* 260/261:589–593
- Sampaio AH, Rogers DJ, Barwell CJ (1998) A galactose-specific lectin from the red alga *Ptilota filicina*. *Phytochemistry* 48:765–769
- Sato Y, Morimoto K, Hirayama M, Hori K (2011) High mannose-specific lectin (KAA-2) from the red alga *Kappaphycus alvarezii* potentially inhibits influenza virus infection in a strain-independent manner. *Biochem Biophys Res Commun* 405:291–296

- Schaefer RL, Keller KF, Doyle RJ (1979) Lectins in diagnostic microbiology: use of wheat germ agglutinin for laboratory identification of *Neisseria gonorrhoeae*. *J Clin Microbiol* 10:669–672
- Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M (2006) In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat Protoc* 1:2856–2860
- Sugahara T, Ohama Y, Fukuda A, Hayashi M, Kawakubo A, Kato K (2001) The cytotoxic effect of *Eucheuma serra* agglutinin (ESA) on cancer cells and its applications to molecular probe for drug delivery system using lipid vesicles. *Cytotechnology* 36:93–99
- Templier V, Roux A, Roupioz Y, Livache T (2016) Ligands for label-free detection of whole bacteria on biosensors: a review. *TrAC Trend Anal Chem* 79:71–79
- Van Stokkum IHM, Spoelder HJW, Bloemendal M, Van Grondelle R, Groen FCA (1990) Estimation of protein secondary structure and error analysis from CD spectra. *Anal Biochem* 191:110–118
- Vasta GR, Ahmed H (eds) (2008) *Animal lectins: a functional review*. CRC Press, Boca Raton
- Whitley MJ, Furey W, Kollipara S, Gronenborn AM (2013) *Burkholderia oklahomensis* agglutinin is a canonical two-domain OAA-family lectin: structures, carbohydrate binding, and anti-HIV activity. *FEBS J* 280:2056–2067
- Whitmore L, Wallace BA (2008) Protein secondary structure analyses from circular dichroism spectroscopy: methods and reference databases. *Biopolymers* 89:392–400